

## INFLUENCE OF PARASITIC INFESTATION ON SERUM PROTEIN AND SOME ENZYMES PROFILE IN SHEEP

Ismail, M.; Hendawy, A.A.\* and Hamada, M.M.Z.\*\*

Biochemistry Dept. Faculty of Vet. Med., Zagazig University

\* Zoology Dept. Faculty of Science, Zagazig University

\*\*Physiology Dept. Faculty of Vet. Med., Zagazig University

### ABSTRACT

This study was designed to investigate the different infestation levels of *Fasciola* on some of the biochemical parameters specially on enzymes and protein using forty three adult male sheep, classified into four groups (control, light, moderate and heavy *Fasciola* infested) according to the intensity of infestation manifested by the fecal egg count. Liver enzymes AST, GGT, and GDH were increased and very diagnostic at all levels of infestation while ALT only increased in moderate infestation. The isoenzyme, fast liver ALP, liver and bone ALP isoenzymes, LDH<sub>4</sub> and LDH<sub>5</sub> were increased. This indicative to the liver damage caused by the migrating flukes. Total serum proteins were also decreased at all levels of infestation with a marked decrease in albumin and A/G ratio accompanied by marked increase in  $\gamma_2$  globulin.  $\gamma_1$  was diagnostic in moderate infestation,  $\beta_1$  in light, and heavy groups while  $\beta_2$  was indicative in moderate and heavy infestation.

### INTRODUCTION

Fascioliasis, is primarily a zoonotic disease that produces liver infection with adult flukes. Fecal examination can be used for diagnosed of *Fasciola* infestation in sheep, but early infestation can not be diagnosed by fecal examination until liver damage associated with the immature migratory stage has occurred and adult flukes are producing ova (Wyckoff and Bradley, 1985). The elevation of certain enzymes may be due to the escape from the disrupted hepatic parenchymal cells or altered membrane permeability e.g. Aspartate aminotransferase (AST), Alanine aminotransferase (ALT) and Lactate dehydrogenase (LDH) or cholestasis e.g. Alkaline phosphatase (ALP) and  $\gamma$ -glutamyl transpeptidase (GGT) Coles (1986). Fascioliasis is known to produce a variety of biochemical effects in the host, including changes in plasma enzyme activities indicative

of liver injury (Lopez, et al., 1993 and Perez et al., 2002).

Isoenzymes of LDH or ALP are useful in diagnosis, because of their variation in properties activities from one tissue to another and determination of the type of isoenzyme presenting the serum in disease, as well as the change in total enzyme activity, can provide valuable additional information Moss and Butterworth (1974). The importance of alkaline phosphates as a diagnostic tool in hepatobiliary disease and in bone disease associated with an increased osteoblastic activity. Different ALP isoenzymes can be introduced into the blood as a result of damage to a number of organs (liver, bone, kidney and intestine). Therefore, an increase in total serum ALP activity is not, itself, a direct reflection of damage to any one organ. For this reason, any attempt to determine the source of an increase in serum ALP activity requires an investigation of its tissue-specific isoenzymes. Because ALP isoenzymes retain their tissue specificity, when released into the circulation, they can be used to identify the tissue contributing the isoenzyme that is responsible for the increase in total serum ALP (Dumas and Spano 1980 and Osamn et al., 1999). In liver diseases, an increase in LDH5 is found, especially acute hepatitis, but also in more chronic conditions such as cirrhosis and obstructive jaundice, even when the total serum LDH activity is within normal range (Elliot and Wilkinson 1963).

GGT is useful clinically in diagnosis of diseases of the liver and biliary tract. Its degree of elevation is only moderate in hepatocellular liver disease, but is marked in obstructive liver disease. GGT helps to determine whether an elevated ALP is coming from bone or liver Kaplan, et al. (1995) and Calleja et al. (2000). Serum Glutamate dehydrogenase (GDH) is a more sensitive indicator of liver necrosis than serum GOT activity Sewell, (1967).

The present work aimed to study the effect of different stages and forms of *Fasciola* infestation on the pattern of Lactate dehydrogenase and Alkaline phosphatase isoenzymes, which may be used as a method of evaluation of the health status of sheep as well as a diagnosis of infestation. Also, studying the enzymatic changes of some liver enzymes, Protein and its fractionation also have been included.

### MATERIAL AND METHODS

A total of 43 male sheep native breed, collected from zagazig and El -Basateen abattoirs, about 20-30 kgm (Body weight) were used in this study. The animals were subjected to pre-slaughtering examination for detection of healthy animals. After slaughtering, blood, fecal and bile samples were collected individually from each animal. Postmortum examination is made on the liver for macroscopic detection of the parasite. Fecal and bile samples were collected from each animal directly from the rectum and the gall bladder in Poly-ethylene nylon sacs. The sam-

ples are transferred directly to the laboratory and examined for detection of fluke eggs according to Kelly (1984) and for determination of the laboratory and examined for according to the method of Coles (1986). The animals are classified into four main groups according to the egg count in both feces and bile:-

Serum samples were used for determination of ALT and AST activities according to the method described by Reitman and Frankel (1957). ALP assayed according to Kind and King (1954). GGT according to Szasz et al. (1969) GDH according to Bergmeyer (1974) and LDH according to the method recommended by the German clinical chemistry society (1970). ALP isoenzyme was estimated according to the methods described by Lee and Kenny (1975) and Chapman, et al. (1987) and LDH isoenzyme by Plumner (1978), Stroeve and Makarova (1989), Manchenko (1994). Total protein concentration by the method of (Richardson 1977) and protein electrophoresis by the methods of Smith (1976), Gowenlock, et. al. (1987) and Hames and Rickwood (1990).

## RESULTS

The study resulted in significant increase in the activity of ALT in moderate infestation group. Significant increases in the activity of AST were observed not only at all degrees of infestation, but also increased according to the intensity of infestation. LDH, GGT, and GDH activities were significantly increased in all groups. ALP activity also increased in the moderate and heavy infestation (Table 1). ALP isoenzymes fraction showed that the fast liver fraction increased with all degrees of infestation; also liver and bone fractions are increased while intestinal fraction did not affected (Table 2). LDH isoenzymes results revealed in a non-significant increase in the activity of LDH<sub>1</sub>, LDH<sub>3</sub> fractions, and LDH<sub>2</sub> fraction increased only in moderate and heavy infestation whereas LDH<sub>4</sub> and LDH<sub>5</sub> fractions were highly significantly increased (Table 3). Decreases in the total protein concentrations in serum of light, moderate, and heavy infested sheep, with decrease in albumin, A/G ratio and increases in globulin (Table 4).

## DISCUSSION

The results of ALT in moderate infestation group may be attributed to the hepatocellular damage caused by the migrating liver flukes as previously recorded by Youssef (1983), Nasser and Abd-Rabo (1994) and Azza (1998). On the contrary, Stromberg, et al. (1985) and Bashandy, et al. (1990) reported non significant elevation in the enzyme level, because of the very low concentration of ALT in the liver of ruminants, which prevents any appreciable leakage of the

enzyme during disease process.

Significant increases in the activity of AST were observed not only at all degrees of infestation, but also increased according to the intensity of infestation. Similar findings were observed by **Radostitis, et al. (1994)** and **Ferre, et al. (1995)** in sheep, and also attributed the changes in AST values to the degree of damage of hepatic cells caused by migrating flukes and the penetration of this enzyme to the blood from cytoplasm and hepatocyte mitochondria. Changes in the enzyme activity allow to determine the extent of damage occurring in the liver parenchyma, which in turn, depends on the intensity of infection.

GGT activity was significantly increased in all groups as recorded by **Jemli, et al. (1993)**, **Ferre, et al. (1994)** and **Ferre, et al. (1998)**.

**Galtier, et al. (1986)** related the initial rise of GGT to the penetration of bile ducts by migrating flukes and would not be released into the circulation until bile duct damage occurs. This was also confirmed by **Sykes, et al. (1980)**, **El-Samani, et al. (1985)** and **Rowlands, et al. (1985)** they reported an elevated level of serum GGT and considered to be indicative of epithelial damage in bile ducts or cholestasis in sheep. The elevated values have been reported with fascioliasis, particularly in the chronic form when adult flukes are present in the bile ducts.

GDH showed a significant increase in all levels of infestation. **Haroun, et al. (1986)** recorded a significant increase in GDH activity in fasciola infested sheep. **Boyd (1962)** reported that serum GDH level was considered to be a high specific indicator for hepatocellular damage and high concentration of GDH was found in ovine liver. **Ferre, et al. (1995)** related the increase in GDH level to the inflammatory state of the liver and to tissue destruction provoked by the parenchymal migration of juvenile flukes during the first stage of fascioliasis.

The results explained in table 2 declared a significant increase in ALP activity especially in the moderate and heavy infestation. These results are in accordance with **Degheldy, et al. (1990)**; **Azza (1998)**; **Cosme et al. (2001)** and **Massoud et al. (2001)**. The significant increase was explained by **Kumar, et al. (1983)** which may be caused by the extra-hepatic biliary occlusion by mature flukes which resulted in ALP excretion through bile and its regurgitation into blood stream. In addition, **Abdou, et al. (1976)** attributed this increase to the irritation of liver cells by toxins or metabolic products of the worm and eggs.

ALP isoenzymes fraction showed that the fast liver fraction increased with all degrees of infestation; also liver and bone fractions are increased while intestinal fraction did not affected. **De-Broe (1985)** explained the origin of the fast liver ALP fraction in sera with cholestasis, its origin

lates from the liver plasma membrane. Bile salts leach the enzyme off the sinusoidal and canalicular membranes. Upon reaching the serum, the membrane bound enzyme aggregates with lipids and lipoproteins. **Collin et al. (1987)** suggested that the measure of intestinal ALP fraction is one method of discriminating intra-hepatic from extra-hepatic jaundice, because intestinal ALP may be absent in extra-hepatic obstruction.

LDH activity in control sheep was  $154.87 \pm 20.04$  u/l which significantly increase in blood of sheep infested with fasciola regardless the degree of infestation light, moderate or severe ( $243.98 \pm 18.45$ ,  $236.10 \pm 20.43$  and  $390.06 \pm 20.41$  u/l respectively. These findings were in agreement with that obtained by **Bhattacharya, et. al. (1992)** who concluded that LDH activity was significantly increased when young flukes migrating through the liver. On the other hand, the activity was reduced when the flukes enter the bile ducts. **Abdel-Razek, et. al. (1988)** attributed the occurrence of high LDH activity to the alternation in cell membrane permeability which may lead to leakage of the enzyme from the cytoplasm to plasma.

Total LDH activity can not consider as an accurate assay for liver status, and isoenzyme determinations are more costly. Results from table 3 revealed a non-significant increase in the activity of LDH1, LDH3 fractions, and LDH2 fraction increased only in moderate and heavy infestation whereas LDH4 and LDH5 fractions were highly significantly increased. These results are in accordance with **Stroev and Makarova (1989)**, **Meyer, et al. (1992)**, **Kaplan, et al. (1995)** and **Teitz (1996)** in human.

**Moss and Butterworth (1974)** remarked that LDH5 isoenzyme which occurs in high concentration in tissues such as liver and skeletal muscles that are able to undertake anaerobic metabolism. So when damage to these organs occur, the release of such isoenzyme of LDH into the blood stream is increased. Thus, comparing the serum isoenzyme pattern with that of the suspected tissue can identify the sources of an elevated LDH. **Schaffner and Schaffner (1987)** said that the activity of LDH isoenzymes were not as great as aminotransferases activity in hepatocellular injury. LDH is the predominant isoenzyme in hepatic anoxia and primary liver disease.

The present work manifested decreases in the total protein concentrations in serum of light, moderate and heavy infested sheep, as previously reported by **Degheidy, et al. (1990)** **Kumar and Sharma (1991)** and **Azza (1998)**. **Furmaga and Gundlach (1967)** referred the decrease of total serum proteins caused by the liver flukes, which seriously affect the protein metabolism through mechanical and toxic injury of the liver cells and changes in the physiological function of the organ. On the contrary **Kadhim (1976)** and **Anderson, et al. (1978)** said that the total protein concentrations were increased, whereas **Agnes and Genchi (1977)** and **Radostits, et al. (1994)** reported no changes in total protein concentrations. **Kadhim (1976)** discussed the

decrease in total serum proteins caused by the migrating juvenile flukes will depend on the size of inoculum and the duration of infection. This phenomenon will coincide with greatest destruction of cells brought about by the juvenile flukes just before they arrive in the bile duct.

The total albumin value was significantly decreased in all infested groups. **Kadhim, (1976) and Waweru et al. (1999)** thought that the hypoalbuminaemia may be due to two factors, one is the decreased synthesis and the other one is the higher catabolic rate due to the damage to the liver parenchyma caused by the migrating juvenile flukes. In addition, **Ghazy (1987)** stated that albumin is lost into the gastrointestinal tract via the bile duct as a result of feeding habits of the flukes.

On dealing with the effect of fasciola on A/G ratio, the data revealed a significant decrease in A/G ratio at all level of infestation. These results are in consistence with that mentioned by **Bashandy, et al. (1990) and Azza (1998)**, which discussed the obtained results for both albumin and globulins .

The non-significant increases in  $\alpha_1$  &  $\alpha_2$  fractions at all levels of infestation revealed. While  $\beta_1$  was significantly decreased in light and heavy infestation,  $\beta_2$  were significantly increased in medium and heavy infestation, while  $\beta_3$  were not significantly changed,  $\gamma_1$  was significantly increased in medium infestation while  $\gamma_2$  was significantly increased at all levels of infestation (Table 4). The investigation of **El- Hawary, et al. (1971)** reported that the presence of parasite could be a stimulatory factor for an increase in the synthesis of gamma globulin which is the main globulin fraction. While **Mulligan (1972)** provided evidence that beta globulin fraction was significantly increased in liver cirrhosis and obstructive jaundice and attributed such elevation to the stimulation of reticulo- endothelial system. **Kadhim (1976)** reported that  $\alpha_1$  globulin is not materially affected by the destruction of hepatic cells or by the presence of parasite.

The use of some hepatic marker enzymes GGT and GDH, which are found to be increased during the early migratory stage of Fasciola infestation. Isoenzyme application also is more useful in determining the origin of the elevated enzyme; also protein electrophoresis is useful indicator for the stage of liver damage. All of these tools can be used as a method of evaluation of the health status of the sheep as well as a diagnosis of infestation and the intensity of infestation.

Groups	No. of sheep	Egg count/gm	Average
Control	10	----	---
Light infestation	11	100-200	150
Moderate infestation	12	200-400	300
Heavy infestation	10	400-800	600

Table (1): represent the mean activity S.E. of ALT, AST, GGT and GDH enzymes in control, light, moderate and heavy *Fasciola* infested sheep (U/l).

Group	ALT	AST	GGT	GDH
Control	22.20± 2.52	130.20±23.98	28.56± 2.21	3.49± 1.35
Light infestation	26.27± 3.08	25.509±22.86*	37.71±2.68*	15.58±3.48*
Moderate infestation	31.67±4.10*	234.33±28.07*	36.79±1.63*	18.20±2.70*
Heavy infestation	29.90± 3.52	243.20±33.20*	42.36±1.28*	17.16±3.53*

\* = Significance

Table(2): ALP isoenzymes in control, light, moderate and heavy *Fasciola* infested sheep (U/l).

Group	Total ALP	Fast liver	Liver & bone	Intestinal
Control	140.02±9.11	00.00±0.0	96.80±6.42	52.22±3.42
Light infestation	206.15±28.21	5.28±0.51*	148.74±21.95	52.12±7.05
Moderate infestation	258.38±28.53*	8.22±1.01*	178.63±21.01*	62.53±5.17
Heavy infestation	256.66±24.73*	13.36±0.72*	197.74±21.14*	45.60±4.6

\* = Significance

Table (3): Discuss LDH isoenzymes in control, light, moderate and heavy fasciola infested sheep (U/l).

Group	Total LDH	LDH <sub>1</sub>	LDH <sub>2</sub>	LDH <sub>3</sub>	LDH <sub>4</sub>	LDH <sub>5</sub>
Control	154.87 ±20.04	83.23 ±11.72	1.29 ±0.19	34.08 ± 5.23	6.93 ± 1.49	26.64 ±3.69
Light infestation	243.98 ±18.45*	81.99 ±6.25	2.06 ±0.20	43.13 ±3.07	24.89 ± 3.83*	91.88 ± 9.70*
Moderate infestation	236.10 ± 20.43*	86.50 ± 7.37	2.48 ± 0.32*	47.40 ± 4.57	25.42 ± 2.83*	74.29 ± 7.83*
Heavy infestation	390.06 ± 20.41	89.79 ± 4.13	2.36 ± 0.42*	42.85 ± 3.58	80.65 ± 5.56*	174.12 ± 11.26*

\* = Significance

Table (4): Interpret the changes in total proteins, and its electrophoretic pattern control, light, moderate and heavy fasciola infested sheep (gm/l).

Group	Control	Light infestation	Moderate infestation	Heavy infestation
Total proteins	70.42± 3.07	58.15± 2.63*	62.54± 1.57*	53.94± 2.35*
Albumin	30.69± 1.30	17.98± 0.93*	17.87± 0.59*	21.67± 0.36*
α <sub>1</sub> Globulin	1.42± 0.010	1.26± 0.08	1.465± 0.07	1.21± 0.08
α <sub>2</sub> Globulin	2.65± 0.16	2.67± 0.15	2.69± 0.15	2.46± 0.15
β <sub>1</sub> Globulin	7.44± 0.37	6.24± 0.37*	7.33± 0.19	5.98± 0.45*
β <sub>2</sub> Globulin	0.81± 0.18	1.02± 0.10	1.48± 0.09*	1.41± 0.25*
β <sub>3</sub> Globulin	6.08± 0.41	6.65± 0.44	9.98± 0.45	5.59± 0.27
γ <sub>1</sub> Globulin	13.75± 0.53	14.84± 0.68	16.52± 0.40*	15.84± 0.94
γ <sub>2</sub> Globulin	7.40± 0.45	9.13± 0.58*	9.43± 0.37*	9.10± 0.45*
A/G Ratio	0.78± 0.02	0.45± 0.01*	0.40± 0.01*	0.32± 0.02*

\* = Significance



### REFERENCES

- Abdel-Razek, M.; El-nagar, W. and Abdel-Raouf, M. (1988) : The significance of serum enzymes and trace elements in addition to fecal egg counts to diagnose and assess pathogenicity of fascioliasis. *Alex.J.Vet.Sci*,4 (1): 581-590.
- Abdou, M. O. (1976) : Clinical studies on liver function tests in buffaloes. M.V.Sc. Thesis. Fac. Vet. Med., Cairo Univ.
- Agnes, F. and Genchi, C. (1977) : Serum enzyme activities in cattle with fascioliasis. *Archivio Veterinario Italiano*.28 (3/4) 131-133.
- Andreson, P. H.; Berret, S. and Patterson, D. S. P. (1978) : Resistance to *Fasciola hepatica* in cattle.2- Biochemical and morphological observation. *J.Comp. Pathol.* 88:245-251.
- Azza, M. Kamal; (1998) : Biochemical changes in bovine serum during parasite infestation. Ph.D. Thesis. Fac. Vet. Med. Cairo Univ.
- Bashandy, M. M.; Safaa Yassein; Lotfy, M. M.; El-Bahy, M.; Amira, M. M. and Dessouky, M. I. (1990) : Hematological and serum biochemical profiles in experimental fascioliasis in sheep. *Egy.J.Com. Pathol. Clin. Pathol.* 3 (2): 357-374.
- Bergmeyer, H. U. (1974) : Methods of Enzymatic Analysis. Vol. 2. Second English edition. Verlag chemie weinheim. Academic press. Inc.
- Bhattacharya, M.; Mukit, A. and Baruah, G. K. (1992) : Dehydrogenases in the liver of cattle naturally infected with *Fasciola gigantica*. *Ind. J. of Ani. Sc.* 62(1): 43- 44.
- Boyd, J. W. (1962) : The comparative activity of some enzymes in sheep, cattle and rat. Normal serum and tissue levels and changes during liver necrosis. *Res. Vet. SC.*, 3: 256-268.
- Calleja, C.; Bigot, K.; Eeckhoutte, C.; Sibille, P.; Boulard, C. and Galtier, P. (2000) : Comparison of hepatic and renal metabolizing enzyme activities in sheep given single or two fold challenge infections with *Fasciola hepatica*. *Int. J. Parasitol.* 30, (8) : 953-958.
- Chapman, J. F. Wooded, L. L. and Silverman, L. M. (1987) : In *Methods in Clinical Chemistry*. By Pesce, A. J. and Kaplan L. A. The C.V. Mosby Comp.
- Coles, H. E. (1986) : *Veterinary Clinical Pathology*. Fourth edition. W.B. Saunders Company, Philadelphia.
- Collin, D.; Goold, M.F. and Rosalki, S. B. (1987) : Plasma intestinal ALP and intermediate molecular mass gamma- glutamyl transferase activities in differential diagnosis of jaundice. *J. Clin. Pathol.* 40 : 1252.

- Cosme, A., Ojeda, E.; Cill, G.; Torrado, J.; Alzate, L.; Beristain, X.; Orive, V. and Arenas, T. (2001)** : Fasciola hepatica. study of a series of 37 patients Gastroenterol hepatol 24(8): 375-80.
- De-Broe, M. E.; Roels, F.; Nouwen, E. J.; Claeys, L. and Wieme, R. J. (1985)** : Liver plasma membrane: The source of high molecular weight ALP in human serum. Hepatology.5 : 118.
- Degheldy, N.S.; Amin, MM and Ghazy, A.A. (1990)**: Effect of fascioliasis on liver function before and after treatment in sheep. Egy. J. Comp. Pathol. Clin. Pathol. 3 (II): 65-76.
- Dumas, M .B.and Spano, J.S. (1980)**: Characterization of equine alkaline phosphatase isoenzymes based on their electrophoretic mobility by polyacrylamide gel disc electrophoresis Am. J. Vets. Res. 41 (12): 2076 -2081.
- El-Hawary, M. F. S.; Ibrahim, A. M.; Shaker, A. H. and Saif, M. (1971)** : Studies on serum proteins in G. I. P. by agar gel electrophoresis and the effect of treatment with nitidazole. Egy. J. Med. Ass., 54:101.
- Elliot, B. A. and Wilkinson, J. H. (1963)** : The serum a hydroxy butarate dehydrogenase in diseases other than myocardial infarction. Clin. Sci. 24 :343.
- El-Samani, F.; Mahmoud, O.M.; Fawi, M. T.; Gameel, A.A. and Haroun, E. M. (1985)** : Some enzyme activity and bilirubin concentration in sheep experimentally infected with Fasciola gigantica. Journal of Comparative Pathology 95 (4): 499-503.
- Ferre, J.; Barrio, J.P.; Gonzalez-Gallego and Vazquez, F.A. (1994)**: Appetite depression in sheep experimentally infected with F.hepatica. Veterinary Parasitology 55:71-79.
- Ferre, L; Lopez, P.; Gonzalo-Orden, M.; Julian, M. D.; Rojo-Vazquez, F. A. and Gonzalez-Gallego, J. (1995)** : The effect of subclinical fascioliasis on hepatic secretory function in sheep. Parasitol.Res. 81: 127-131.
- Ferre, L; Lopez, P.; Vazquez, F.A. and Gallego, J.G. (1996)** : Experimental ovine fascioliasis: Antipyrine clearance as indicator of liver damage. Veterinary Parasitology 62: 93-100.
- Furmaga, S. and Gundlach, J. L. (1967)** : Studies on the serum proteins in experimental fascioliasis in sheep. Acta Para. Polonica. XI 5 /30 (11) Fasc. 13.
- Galtier, P. Larrieu, G. Tufenkji, A. and Franc, M. (1986)**: Incidence of Experimental fascioliasis on the activity of drug-metabolising enzymes in lamb liver Drug metabolism and Disposition 14,137-141.
- García, L. S. and Bruckner, A. D. (1993)** : Diagnostic Medical Parasitology. Second edition.

American Society for Microbiology. Washington, D.C.

- Ghazy, A. A. (1987)** : Some studies on fascioliasis in sheep in Egypt M. V.Sc., Fac. of Vet. Med., Cairo Univ.
- Gowenlock, A. H.; Memurray, J. R. and Mclauchlan, D. M. (1987)** : Varley's Practical Clinical Biochemistry. Sixth edition. Heinemann Medical Books, London.
- Hames, H. D. and Rickwood, D. (1990)** : Gel Electrophoresis of Protein. A Practical Approach. Second edition. Orl Press at Oxford University Press.
- Haroun, E. M.; Gadire, H. A. and Gameel, A. A. (1986)** : Studies on naturally occurring ovine Fascioliasis in Sudan J. Helminth. 60 (1): 46-53.
- Jemli, M. H.; Braun J. P.; Darchies, P.; Romdhane, M. N. and Kilani, M. (1993)** : Biochemical and hematological studies on lambs infected experimentally with *F. hepatica*. *Revue de Medecine Veterinaire* 196 (4): 241-249.
- Kadhim, J. K. (1976)**: Changes in the serum protein values of sheep infected with *F. gigantica*. *Am. J. Vet. Res.* 37 (2): 229-231.
- Kaplan, A.; Jack, R.; Ophelm, K.; Toivola, B. and Lyon, A.W. (1995)**: Clinical Chemistry :- Interpretation And Techniques. J Fourth edition. Williams and Wilkins.
- Kelly, W. R. (1984)**: Veterinary Clinical diagnosis 3rd Ed. Balliere. Tindall New York
- Kind, P.R.N. and King, E.G. (1954)**: Estimation of plasma phosphatase by determination of hydrolyzed phenol with amino-antipyrine. *J. Clin. Path.* 7: 322.
- Kumar, M.; Maru, A. and Pachauri, S. P. (1983)**: Role of enzymes in evaluating flukicidal action of Disophenol in fasciola affected buffaloes. *Br. Vet. J.* 193: 262.
- Kumar, S. and Sharma, M. C. (1991)**: Infertility of rural cows in relation to fascioliasis. *Indian J. Ani. Sci.* 61 (8) 838-840.
- Lee, L. M. Y. and Kenny, M. A. (1975)**: Electrophoretic methods for assessing the normal and pathological distribution of ALP isoenzymes in serum. *Clin. Chem.* 21(8): 1128-1135.
- Lopez, P.; Tunon, M. J.; Gonzalez, P.; Diez, N.; Bravo, A. M. and Gonzalez-Gallego, J. (1993)**: Ductular proliferation and hepatic secretory function in experimental fascioliasis. *Exp. Parasitol.* 77 (1): 36-42.
- Manchenko, G. P. (1994)**: Hand Book Of Detection Of Enzymes On Electrophoretic Gels. CRC press.
- Massoud, A.; El-Sisi, S.; Sulama, O. and Mussoud, A. (2001)**: Preliminary study of therapeutic

efficacy of a new fasciolicidal drug derived from commiphora molomol (Myrrh) Am. J. Trop. Med. Hyg. 65 (2): 96-9.

**Meyer, D.I.; Coles, E. H. and Rich, L.J. (1992):** Veterinary Laboratory Medicine. Interpretation and Diagnosis First edition. W.B. Saunders Company. Philadelphia.

**Moss, D.W. and Butterworth, P.J. (1974):** Enzymology And Medicine. First edition. Pitman Medical Publishing Company LTD.

**Mulligan, W. (1972):** The effect of helminthic infection on the protein metabolism of the host. Proc. Nut. Soc., 31: 47-51.

**Nasser, M.H. and Abd Rabo, T. M.A. (1994):** The efficacy of fastnex against natural infection of cattle by fascioliasis in Kahr El-Sheikh Governorate. Assiut Vet. Med. J. 31(62): 229-245.

**Osman, M.M.; Ismail, X. and Aref, T.Y. (1999):** Human fasciolosis : a study on the relation of infection intensity and treatment of hepatofiliary affection. J. Egypt Soc. Parasitol. 29 (2): 353-363.

**Perez, J.; Ortega, E.; Moreno, T.; Morrondo, P.; Lopez-Sandoz, C. and Matti, Nez-Moreno, A. (2002):** Pathological and immunohistochemical study of the liver and hepatic lymph nodes of sheep chronically reinfected with *Fasciola hepatica*, with or without Triclabendazole treatment. J. Comp. Pathology. 127 (1): 30.

**Plummer, D.T. (1978):** An Introduction to Practical Biochemistry. Second edition McGraw-Hill Books Company LTD.

**Radostits, O.M.; Blood, D.C. and Gay, C.C. (1994):** Veterinary Medicine. Eighth edition. Balliere Tindall.

**Reitman, S. and Frankel, S. (1957):** A calorimetric method for determination of serum glutamic oxalic transaminase and glutamic pyrovic transaminase. Am. J. Clin. Path. 28 : 56.

**Richardson, T. (1977):** Annals of Clinical Biochemistry, 14: 223-226. Quoted in Medical Laboratory Manual for Tropical Countries By Monica, C. (1987) Vol.1, Second edition. The Univ. Press., Cambridge, G.B.

**Rowlands, D. Ap. T.; Clampitt, R.B. and Macpherson, I.S. (1985):** The ability of dioamphenithide to control immature *Fasciola hepatica* in sheep at a lower than standard dose level. Vet. Rec. 116: 182-184.

**Schaffner, J. A. and Schaffner, E. (1987):** Quoted in Methods in Clinical Chemistry. By Pesce, A. J. and Kaplan L.A. The C.V. Mosby Comp.

- Sewell, M.M.M. (1967):** Serum enzyme activities in acute ovine fascioliasis. *Vet. Record* 80 (19): 577-578.
- Smith, I. (1976):** *Chromatographic and Electrophoretic Techniques*. Vol. 2. Zone Electrophoresis Fourth edition. William Heinemann Medical Books LTD. London.
- Stroev, E. A. and Makarova, V. G. (1989):** *Laboratory Manual In Biochemistry*. Mir Publishers. Moscow.
- Stromberg, B. E.; Conboy, G.A.; Hoyden, D.W. and Schlotthaue, J.C. (1985):** Pathophysiological effects of experimentally induced *Fasciola magna* infection in sheep. *American Journal of Veterinary Research* 46 (8): 1637-1641.
- Sykes, A. R.; Coop, R.L. and Robinson, M. B. (1980):** Chronic subclinical ovine fascioliasis : Plasma Glutamate dehydrogenase, Gamma glutamyl transferase and Aspartate aminotransferase activities and their significance as diagnosed aids. *Res.Vet. Sci.* 28 (1): 71-75.
- Szasz, G.; Rosenthal, P. and Fritzsche, W. (1969):** Die glutamyl transpeptidase aktivitat in serum bei hepato-biliaren Erkrankungen. *Deutsche Med. Wochenschrift* 94 : 1911.
- Tietz, N. W. (1996):** *Fundamental of Clinical Chemistry*. Forth edition. Vol. 1. (Moss, D. W. and Henderson, A.R.) W.B. Sanders Company P.P. 283-350.
- Waweru, J.G.; Kanxari, P.W.; Mwangi, D.M.; Nagtia, T.A. and Nansen, P. (1999):** A comparison of serum biochemical changes in two breeds of sheep (Red Masai and Dorper) experimentally infected with *Fasciola gigantica*. *Onderstepoort J. Vet. Res.* 66(1): 47-49.
- Wyckoff, J. H. and Bradley, R. E. (1985):** Diagnosis of *Fasciola hepatica* infection in beef calves by plasma enzyme analysis. *Am. J. Vet. Res.*, 46(5): 1015-1019.
- Youssef, H. F. (1983):** Study of liver fluke infection in cattle with special reference to methods of diagnosis and treatment together with changes in constituents and minerals. Ph.D. V.Sc. Thesis., Fac. Vet. Med. Zag. Univ.

الملخص العربى

تأثير الإصابة بالديدان على بروتينات الدم  
وبعض الإنزيمات فى الأغنام

محمد إسماعيل محمد      أحمد هندأوى\*      محمد زكى حماده\*\*

قسم الكيمياء الحبرية - كلية الطب البيطرى - جامعة الزقازيق

قسم الفسيولوجيا\* - كلية الطب البيطرى - جامعة الزقازيق

قسم الحيران - كلية العلوم - جامعة الزقازيق\*\*

أجريت هذه الدراسة على ثلاثة وأربعون من ذكور الأغنام تم تقسيمها إلى أربعة مجموعات (مجموعة ضابطة ومجموعة بسيطة الإصابة ومجموعة شديدة الإصابة) على أساس عدد البيض فى البراز زيادة إنزيمات الكبد الأسبارتات أمينوترانسفيراز وجاما جلوتاميل ترانساميناز وجلوتامات ديهيدروجيناز فى كل درجات الإصابة تعتبر من وسائل التشخيص بينما زاد إنزيم الألانين أمينوترانسفيراز فى المجموعة متوسطة الإصابة فقط أشباه الإنزيمات (جزء الكبد السريع وجزء الكبد والعظم لإنزيم الفوسفاتيز القلوى وكذلك الجزء الرابع والخامس من إنزيم اللاكتات ديهيدروجيناز) تدل على تأثر الكبد بالديدان المهاجرة داخل خلاياها. قلة مستوى البروتينات الكلية فى جميع درجات الإصابة مع قلة مستوى الزلال ونسبة الألبومين إلى الجلوبيولين.