

- 16. Which of the following disorders demonstrates a mitochondrial inheritance pattern?**
- Retinoblastoma
 - Leber hereditary optic neuropathy
 - Retinitis pigmentosa
 - Fabry disease
- 17. Episcleritis**
- Should be treated with topical corticosteroid or anti inflammatory agents
 - Usually is associated with a systemic disease that should be treated
 - Causes severe ocular pain
 - Is almost always a self-limited condition that, if untreated, runs its course in a few days
- 18. The second most common location for basal cell carcinoma is:**
- Upper lid
 - Lower lid
 - Medial canthus
 - Outer canthus
- 19. True statement regarding posterior subcapsular cataracts include**
- Blurring is greater in near than far vision
 - Blurring is greater at night than day light
 - They are often the sequelae of penetrating lens injury
 - One cause is the use of amiodarone
- 20. Earliest sign of pilloedema is:**
- Obliteration of optic cup
 - Loss of venous pulsation
 - Blurring of disc margins
 - Hyperaemia of the disc
- 21. After thorough and copious irrigation of conjunctival fornices, the next step in initial management of a patient with a chemical burn:**
- Topical steroid agents
 - Topical ascorbate
 - Topical antibiotic agents
 - Debridement of any foreign particles
- 22. Substance deposited in band shaped keratopathy is?**
- Calcium phosphate
 - Magnesium phosphate
 - Magnesium sulphate
 - Iron
- 23. Most common second malignancy in survivors of retinoblastoma is?**
- Optic glioma
 - Thyroid cancer
 - Papillary thyroid tumor
 - Osteosarcoma