

## **POPULATION FLUCTUATIONS OF SOME INSECT PESTS INFESTING BROAD BEAN PLANTATIONS IN RELATION TO CERTAIN ECOLOGICAL FACTORS**

**Shalaby, H. H. ; E. M. A. Mousa and Samia A. El-Gawwad**  
Plant Protection Res. Institute, ARC, Dokki- Giza, Egypt

### **ABSTRACT**

Field experiments were conducted at Kafr El-Sheikh governorate during two successive seasons 2008/2009 and 2009/2010 to study the population fluctuations of some insect pests infesting broad bean plantations namely *Liriomyza trifolii* (Burgess) ; *Aphis craccivora* (koch) and *Empoasca discipiens* Poali. In addition, the effect of certain weather factors (daily mean temperatures and daily mean R.H.) and plant age were studied on the population fluctuations of the previously mentioned pests. The weather factors and plant age had significant effect on the population fluctuations of *L. trifolii*, *A. craccivora* and *E. discipiens*. Also, the relative humidity had shown no significant effect on population fluctuations of the three insect pests during the two seasons 2008/2009 and 2009/2010. The percentage of Explained variance was 83.5 & 81.9 % during the two seasons, respectively. The phytochemical contents of the broad bean plant stages, in leaves of Foliage phase, of *A. craccivora* ; *L. trifolii*, and *E. discipiens* were significant during two successive seasons.

### **INTRODUCTION**

Broad bean, (*Vicia faba* L) is becoming one of the most important vegetable crops grown for local consumption. *Liriomyza trifolii* (Burg.) (Diptera: Agromyzidae) was recorded on cotton in some Governorates in Egypt. Chang and Chen (1993) concluded that *L. trifolii* is a potential pest in the cotton fields in Egypt and more research is needed (Abd-El-Wahab ;1998 and Shalaby *et al* 2009. Throughout the growing season, broad bean plants are liable to infestation by phytophagous pests such as *Aphis craccivora* (koch), *Empoasca discipiens* Poali, which considered the most common and important insect pests of broad bean plants. In heavy infestation, these pests are causing serious damage to plants, leading to great reduction in the yield (Helal *et al* 1996 and El-Defrawi *et al.* 2000). Aphids play an important role as a vector of plant viruses and produce honeydew (Li *et al* 1994). Therefore, the purpose of this work was to study the effect of plant age and certain weather factors on the population fluctuations of some insect pests infesting broad bean and the effect of insect infestation on chemical components of broad bean leaves.

### **MATERIALS AND METHODS**

Experiments were carried out at Plant Protection Institute Experimental at Kafr El-Sheikh Governorate during two successive seasons, (2008/2009 and 2009/2010) . An area of one feddan was sown with broad bean, seeds (*Vicia faba* L) variety ; Balady on November 10<sup>th</sup> & 15<sup>th</sup> during 2008 and 2009.

Population study was conducted in an area of one feddan, divided into four replicates. The plants received all normal recommended agricultural practices of bean seeds with the absence of any insecticides application. Population study of *Liriomyza trifolii*, (Burg.); *Aphis craccivora* (koch) and *Empoasca discipiens* Poali started after about one month from sowing and continued until the end of the season.

Weekly direct count of the gassid, *Empoasca discipiens* Poali was done by counting its numbers on sample of 80 leaves chosen randomly (20 leaves/replicate) Four replicates were chosen randomly early in the morning before the gassid adults tend to be more active (Gameel, 1973).

After direct count of adults, the previous samples were picked and then put in a paper bags and examined on the same day in the laboratory with the aid of a stereoscopic-microscope for counting the legume aphid, *Aphis craccivora* (Koch) "nymphs and adults" the broad bean leaf miner, *Liriomyza trifolii* (Burgess) "larvae".

The records of meteorological data, the daily mean of minimum, maximum temperature and daily mean relative humidity were obtained from the meteorological records of Central Laboratory for Agriculture Climate, Agriculture Research Center at Dokki, The daily records of these factors were recalculated to get the daily averages within one week of the sampling date.

#### **Chemical analysis of broad bean leaves:**

The samples were cleaned and washed with distilled water. The fresh weight was determined, and then leaves were put in a drying oven at 60°C for one day. The dry leaves was stored in glass bottles to devote for total carbohydrate, total sugar (reduced sugar and non-reduced sugar), total nitrogen and total lipid.

Total nitrogen was determined according to the method of micro kjeldahl as described by Peach and Tracy (1956). Total carbohydrates were determined according to Forsee (1941). Readings were taken calorimetrically using spectromic 20 with 420 mm wave length. It was calculated according to the following equation:

$$\frac{C/n = \text{carbohydrate content (gm)}}{\text{Nitrogen content (gm)}}$$

#### **Statistical analysis:**

The statistical analysis (ANOVA and Simple correlation) of the obtained data were performed by using SAS program (SAS Institute, 1988) which run under WIN.

## **RESULTS AND DISCUSSION**

Population fluctuations studies showed that *Liriomyza trifolii* (Burgess); *Aphis craccivora* (koch)., and *Empoasca discipiens* Poali, were studied during two seasons (2008/2009 and 2009/2010).

The data in Table (1) show that the infestation of broad bean with *L. trifolii* were started at 29 days after sowing on 9<sup>th</sup> December (7 individuals

/20 leaves) then it was increased sharply to reach its maximum (85 individuals /20 leaves) at 57 days after planting date on 7<sup>th</sup> January. After that the insect population fluctuated and decreased gradually to moderate level (39 individuals /20 leaves) on 21<sup>st</sup> January after 71 days from planting and fluctuated and increased to reach (291 individuals/ 20 leaves) at 11<sup>th</sup> February. Finally the insect population decreased sharply to reach a lower level (1 individual/ 20 leaves) at 141 days after planting on 2<sup>nd</sup> April. These results were in agreement with those of Sharaf El-Din *et al.* (1997).

The first appearance of the broad bean aphid *A. craccivora* was at 22 days after sowing on 2<sup>nd</sup> December (26 individuals /20 leaves) then it was increased sharply to reach its maximum (73 individuals /20 leaves) at 57 days after planting date on 7<sup>th</sup> January. After that the insect population fluctuated and decreased gradually to reach (5 individuals /20 leaves) on 21<sup>st</sup> January after 92 days. It is that population of *A. craccivora* was obviously higher on young plants than on older one. These results are in agreement with findings of Li *et al* and 1994 Helal *et al.* 1996 .

The results indicated that the infestation of the broad bean leaves with *E. discipiens* appeared after 29 days from planting date on 9<sup>th</sup> December 2008(1 individuals/20 leaves) then disappeared absolutely until 7 January 2009 (16 individuals/ 20 leaves) after 57 days from planting. After that the numbers were increased sharply after 85 days on 4<sup>th</sup> February and reached 57 individuals/ 20 leaves . Finally, the insect population decreased gradually on 25<sup>th</sup> March after 134 days from planting date (7 individuals/ 20 leaves). These results indicated that high number of insects infested the moderate plants of broad bean than the younger one , (Table 1).

In Table (2), the results indicated insignificant correlation between the populations of insects and mean daily maximum, minimum temperatures ( $r=0.511$  &  $0.793$ ), during 2008/2009 season. In addition, significant correlation was found between the insect populations and daily relative humidity ( $r=0.774$ ) during the season of 2008/2009 . The partial regression analysis for the effect of weather factors on the populations revealed that means of daily maximum and minimum temperatures had significant effect. during the season of 2008/2009. Also, the means of daily relative humidity had significant effect. These results are in agreement with El- Khouly *et al.* (1998).

The obtained results revealed that the combined effect of the tested plant age and weather factors was significant on the insect population where the calculated "F" value was 11.60.

The analysis of the variance revealed that the weather factors and the plant age are responsible for about 83.5% of the variability in the populations of the observed pests on the broad bean during 2008/2009 season.

In Table (1) show that broad bean leaves infestation with *L. trifolii* were started at 43 days after sowing on 28<sup>th</sup> December (2.33 individuals /20 leaves) then it was increased sharply to reach its maximum (123 individuals /20 leaves) at 85 days after planting date on 9<sup>th</sup> February. After that the insect population fluctuated and decreased gradually to moderate level (72 individuals /20 leaves) on 16<sup>th</sup> February after 92 days from planting and

fluctuated and increased to reach (184 individuals/ 20 leaves) at 2<sup>nd</sup> March. Finally the insect population fluctuation decreased sharply to reach a lower level (12.33 individuals/ 20 leaves) at 141 days after planting on 7<sup>th</sup> April.

The broad bean leaves infested with *A. craccivora* after 22 days of planting on 17<sup>th</sup> December 2009 (5 individuals/ 20 leaves). Then the population increased to reach 11 individuals/ 20 leaves after 36 days of planting on 21<sup>st</sup> December 2009. After that the insect population decreased to reach 1 individual/ 20 leaves after 57 days of planting on 12<sup>th</sup> January 2009. These results indicated that *A. craccivora* preferred the younger globe bean plants than the older ones. These results are in agreement with Shalaby (2004) and Shaalan. (2005).

Data in Table (1) showed that the infestation of broad bean leaves with *E. discipiens* started on 7<sup>th</sup> December 2009 after 36 days of planting date (7 individuals/ 20 leaves). Then, the population increased rapidly to reach 33 individuals/20 leaves after 85 days from cultivation on 9<sup>th</sup> February. Finally, the populations decreased. These results indicated that *E. discipiens* preferred the older broad bean plants than the younger one.

**Table (1) : Population fluctuation of some insects on broad bean plants at Kafr El-Sheikh Governorate during 2008 / 2009 and 2009 / 2010 seasons.**

Inspections after plantation (Week)	Plant age (days)	2008/2009			Inspections after plantation (Week)	Plant age (days)	2009/2010		
		Leaf miner Larvae	Aphid	Jassid			Leaf miner Larvae	Aphid	Jassid
2/12	22	0.00	26.0	0.00	7/12	22	0.00	5.00	0.00
9/12	29	7.00	28.0	1.00	14/12	29	0.00	13.00	0.00
16/12	36	63.00	72.0	0.00	21/12	36	0.00	11.00	7.00
23/12	43	43.00	63.0	0.00	28/12	43	2.33	8.00	4.00
30/12	50	57.00	90.0	0.00	5/1	50	3.67	3.00	5.00
7/1	57	85.00	73.0	16.00	12/1	57	9.00	1.00	7.00
14/1	64	63.00	23.0	4.00	19/1	64	24.67	0.00	8.00
21/1	71	39.00	5.00	8.00	26/1	71	33.67	0.00	2.00
28/1	78	95.00	0.00	4.00	2/2	78	46.33	0.00	4.00
4/2	85	79.00	0.00	57.00	9/2	85	123.0	0.00	33.00
11/2	92	291.00	0.00	4.00	16/2	92	72.00	0.00	12.00
18/2	99	159.00	0.00	18.00	23/2	99	108.0	0.00	31.00
25/2	106	130.00	0.00	19.00	2/3	106	184.67	0.00	12.00
4/3	113	90.00	0.00	17.00	9/3	113	150.67	0.00	15.00
11/3	120	36.00	0.00	17.00	16/3	120	63.00	0.00	16.00
18/3	127	3.00	0.00	18.00	23/3	127	36.33	0.00	9.00
25/3	134	4.00	0.00	7.00	30/3	134	25.33	0.00	14.00
2/4	141	1.00	0.00	14.00	7/4	141	12.33	0.00	3.00
Total	-	1245.00	380.	204.0	Total	-	895.0	41.00	182.00
Mean	-	69.17	21.11	11.33	Mean	-	49.72*	5.00	10.11
LSD			9.6		LSD			5.2	

\*= Significant at 5% level.      A    LSD=1.7                      LSD=6.0                      LSD=Non-significant



the populations revealed that means of daily maximum, minimum temperatures and daily relative humidity had significant effect during the season of 2009/2010. These results are in agreement with El- Khouly et al. (1998).

The obtained results revealed that the variance of combined effect of the tested plant age and weather factors was 81.9% on the insects population where the calculated "F" value was 8.75.

Generally, the results in Table (3) of the two seasons of 2008/2009 and 2009/2010 showed that incidence of infestation with *L. trifolii* increased gradually with the increase of plant age during the two seasons. This result indicated clearly fruiting phase plants are more suitable for *L. trifolii*.

*A. craccivora* during the first season, and second season's infestation occurred on the stages foliage only in both seasons.

*E. discipiens* during two seasons increased gradually with the increase of plant age. This result indicated clearly fruiting phase plants are more suitable for *E. discipiens*.

**Table (3): Rate of infestation of broad bean by some insects and, plant stage during 2008 / 2009 and 2009 / 2010 seasons.**

Insects	Plant age	1 <sup>st</sup> Season:2008/2009		Plant age	2 <sup>nd</sup> Season:2009/2010		Average two seasons
		Plant stages.	Infested plant		Plant stage	Infested plant	
Leaf miner	78	Foliage phase	456	78	Foliage phase	119.7	287.9
	92	Blooming phase	370	92	Blooming phase	195	282.5
	99-141	fruiting phase	423	99-141	fruiting phase	580.3	501.7
Aphids	78	Foliage phase	380	78	Foliage phase	41	210.5
	92	Blooming phase	0	92	Blooming phase	0	0.0
	99-141	fruiting phase	0	99-141	fruiting phase	0	0.0
Jasid	78	Foliage phase	33	78	Foliage phase	37	35
	92	Blooming phase	61	92	Blooming phase	45	53
	99-141	fruiting phase	110	99-141	fruiting phase	100	105

The relation between phytochemical components of broad bean and plant stages infested of *A craccivora* ; *L. trifolii*, and *E. discipiens* are shown in Table (4) indicated that the simple correlation between the phytochemical contents of the broad bean plant stages, in leaves of foliage phase, infested with of *A. craccivora* ; *L. trifolii*, and *E. discipiens* were significant during two seasons.

A significant correlation was found between the two insects, *L. trifolii*, and *E. discipiens* and the content in the blooming phase and fruiting phase while no infestation with *A. craccivora* in two plants stages.

The relation between the phytochemical contents of protein, carbohydrate and lipids contents in broad bean were significantly differed (P<0.01, ANOVA, Duncan's multiple range test). Where, the infested leaves

have the higher contents of Protein, carbohydrate, ratio indicated low ratio in infested leaves.

In this regard, the authors from their opinion stated that, the relationship between phytochemical contents of broad bean leaves and the infestation of *A craccivora* need more investigation in Foliage phase ; infestation of *L. trifolii* and *E. discipiens* investigation in Foliage phase; the blooming phase and fruiting phase, These results agree with those of Helal *et al.* (1996) , Abd-El-Wahab (1998) and Shalaby *et al.* (2009).

## REFERENCES

- Abd El-Wahab, H. (1998). Susceptibility of five broad bean cultivars to the infestation by *Bemisia tabaci* (Gennadius) (Aleyrodidae) *Aphis fabae* (Scop.) (Aphididae) and *Lirimyza trifolii* (Burg.) (Agromyzidae). Egypt J. Appl. Sci., 13 (3): 288-295.
- Chang, D.C. and C.C. Chen (1993). Population fluctuation of major insect pests on kidney bean and the proper time of control. Bull. of Taichung District Agricultural Improvement Station, 38: 11-22.
- El-Defrawi, G. M.; A. K. Emam; L.A. Marzouk and L. Rizkalla, (2000). Population dynamics and seasonal distribution of *Aphis craccivora* Koch and associated natural enemies in relation to virus disease incidence in faba bean fields. Egyptian J. Agric. Res., 78(2): 627-641.
- El-Khouly, A.S.; E.M.E. Khalafalla; M.M. Metwally, H.M. Helal and A.B. Mezaïen (1998). Seasonal abundance and population dynamics of certain sucking insects on Soya bean in Kafr-El-sheikh Governorate, Egypt. J. Agric. Res., 76(1): 141-151.
- Forsee, W.T. (1941): Determination of sugars in plant materials, a photochlorimetric method. Indas. Eng. Chem. Anal. 10<sup>th</sup> ed.: 411 – 418.
- Gameel, O. I. (1973): Field evaluation of insecticides for jassid, *Empoasca lybica* De Berg and whitefly, *Bemisia tabaci* (Genn.) control on cotton. Bull. Entomo. Soc. Egypt, Econo. Ser., 7: 113-122.
- Helal, H.A.; R.M. Salem, A.S. El-Khouly; M.M. Metwally and A.B. El-Mezaïen (1996). Population dynamics of *Aphis craccivora* (Koch) and *Empoasca* spp. on faba bean in relation to associated predators and some climatic factors. Egypt. J. Agric Res. 75 (2) : 461 – 471.
- Li, D. X.; J. Tiang, and Z.Y. Zhang (1994). A study on the bionomics and control techniques of *Aphis craccivora* Koch. on mung beans. Plant Protection, 20(6): 4-5.
- Peach, K. and Tracy, M.V. (1956). Modern methods of plant analysis Springer Vailage. Berlin 1: 479 – 481.
- Shalan, H.S. (2005): Studies on some soyabean pests and their control with non traditional methods. Ph.D.Thesis Fac. of Agric. AL-Azhar Univ. 287 pp.
- Shalaby, S. H.(2004): Studies on the efficiency of some new pest control measures against certain pests of common bean. Ph. D. thesis, Fac. Agric. Moshtohor, Zagazig Univ., 265 pp.

Shalaby, H.H.; Abd-El-Wahab, A.H. and Samia, A.A.Y. (2009). Assessment of *Liriomyza trifolii* (Burg.), (Diptera: Agromyzidae) in cotton (*Gossypium barbadense*) fields in different governorates. Bull. Ent. Soc. Egypt. 45-54

Sharaf El-Din, N.; H. Kannan; A. Babiker; S. Weigand and M.C. Saxena (1997): The leaf miner, *Liriomyza trifolii* Burgess (Diptera: Agromyzidae) a serious insect pest of faba bean (*Vicia faba* L.) in the Sudan. 1. Identification of the pest, natural enemies and abundance. Arab J. Plant Protection. 15(1): 39-42.

### ديناميكية التعداد لبعض الآفات الحشرية التي تصيب زراعات الفول وعلاقتها ببعض العوامل البيئية

حسن شلبي شلبي ، عصام على محمد موسى و سامية عبد الجواد  
معهد بحوث وقاية النباتات- مركز البحوث الزراعية- الدقى- الجيزة

أجريت التجارب الحقلية في محافظة كفر الشيخ خلال سنتين متتاليتين ٢٠٠٨ / ٢٠٠٩ ،  
٢٠١٠ / ٢٠٠٩ لدراسة ديناميكية التعداد لبعض الآفات التي تصيب زراعات الفول وهي حشرة منّ  
البقوليات ، *Aphis craccivora* Koch ، نطاط أوراق البطاطس (*Empoasca*  
*discipiens* Poali و ذبابة أوراق الفول *Liriomyza trifolii* Burgess) وأيضا دراسة  
تأثير بعض العوامل الجوية وهي درجة الحرارة الصغرى والكبرى والرطوبة النسبية وعمر النبات  
على التغيرات في تعداد الآفات المذكورة .

وأظهرت التحليلات الاحصائية للعلاقة بين اعداد الحشرات وبعض العوامل البيئية  
(الحرارة العظمى والصغرى والرطوبة النسبية وعمر النبات) وجود علاقة ارتباط معنوية مختلفة  
بين درجة الحرارة العظمى والصغرى وتعداد الحشرات محل الدراسة ، بينما بالنسبة للرطوبة  
النسبية. توجد علاقة ارتباط معنوية مختلفة علي نبات الفول خلال عامي الدراسة وكانت نسبة  
الاختلاف بين العوامل ٨٣.٥ ، ٨١.٩ % خلال عامي الدراسة.

قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة  
مركز البحوث الزراعيه

أ.د / عادل حسن عبد السلام  
أ.د / حورية عبد الوهاب



