

**THE PREDACEOUS EFFICIENCY OF *Labidura riparia* PALLAS REARED ON RED PALM WEEVIL, *Rhynchophorus ferrugineus* (OLIVIER), IMMATURE STAGES (COLEOPTERA: CURCULIONIDAE)**

**Abdel-Salam, A. H.<sup>1</sup>; A.A.EL-Bana<sup>2</sup>; and Eman E. H. El-Rehewy<sup>3</sup>**

**1 Economic Entomology Department, Faculty of Agriculture, Mansoura University, Mansoura, Egypt. E-mail: adhabdelus@gmail.com**

**2 Chief Researcher and Director of Date palm Research Laboratory- Vice President of Agricultural Research Center, Ministry of Agriculture, Giza, Egypt.**

**3 Plant Protection Research Institute, Agricultural Research Center, Ministry of Agriculture, Dokki, Giza, Egypt.**

**E-mail: [emanelrehewy@yahoo.com](mailto:emanelrehewy@yahoo.com)**

**ABSTRACT**

Laboratory experiments were carried out in order to evaluate the predaceous efficiency of the earwig, *Labidura riparia* Pallas against the red palm weevil (RPW), *Rhynchophorus ferrugineus* immature stages. Obtained results showed that the 6<sup>th</sup> nymphal instar of *L. riparia* was the most predacious stage on all RPW immature stages. However, the 1<sup>st</sup> nymphal stage of *L. riparia* had the least predacious rate on all RPW immature stages. In addition, results showed that all immature stages of RPW were preferred for consumption by the earwig nymphs. The rate of consumption of immature stages of the RPW was increased by the age of the nymphal stage. Results showed that the 6<sup>th</sup> instar nymphs had the longest feeding duration upon all immature stages of RPW. Finally, the earwig, *L. riparia*, can be considered as a promising natural enemy to minimize the red palm weevil population density.

**Keywords:** Red palm weevil, *Labidura riparia*, laboratory experiments, predaceous efficiency.

**INTRODUCTION**

The date palm and the date fruits are hosts for many insects and diseases which are seriously enough to inflict heavy losses if left without control. Under traditional date palm culture, the growers were helpless and in some cases they were unable to identify the causal organism. Date palm insect pests, in general, and the red palm weevil (RPW), *Rhynchophorus ferrugineus* (Olivier) (Coleoptera: Curculionidae), in particular, are widely accepted as being the most destructive factors of date, coconut and oil palms throughout South and Southeast Asia (Wattanapongsiri, 1966). Nowadays, the date palm crop in Eastern Arab countries is under threat. Red palm weevil was probably introduced to the Middle East on infested ornamental palm from India or Pakistan. Red palm weevil was firstly discovered attacking palm in the Arabian peninsula especially United Emirates at 1986 and progressively spread to Gulf states and crossed the red sea into North Africa as the latest record since 1992 in Egypt. It is found over a wide geographical area in Asia, involving many different agro ecosystems. The related species is highly polyphagous with number of known hosts exceeding more than ten different palm species (Murphy and Briscoe, 1999). Effective methods for control this

pest has been difficult to develop. During the last two decades all efforts to control *R. ferrugineus* in the Arab countries, focused on the use of traditional insecticides, modified cultural practices and recently pheromone traps (Abraham *et al.*, 1998). Despite the intensive efforts and the high costs of controlling RPW, it is continuously spreading everywhere and destroying the holy date palm tree. Management programs of RPW depend mainly on chemical insecticides (Girgis *et al.*, 2002). About 900 species of earwigs are identified worldwide, most of them being tropical. Earwigs are active predators of tiny insects, besides feed on living and dead plants of the same ecosystem. They are considered important predators of eggs, larvae and pupae of moths and beetles (El-Hussieni, 1969). The Indian earwig, *Labidura riparia* Pallas, is the dominant species and mainly preys cutworms, caterpillars, small grubs, maggots, mealybugs, fleas and other insects (Dugard, 2006). In Brazil, *Euborellia annulipes* (Lucas) and *L. riparia* were associated with commercial poultry ranches as biological control agents (Guimaraes *et al.*, 1992). In the USA, *L. riparia*, was recorded as a predator of pupal stage of lepidopterous pests in soybean fields (Domiciano and Herzog, 1990 and Lee *et al.*, 1990), and was the most abundant predator in peanut fields (Kharboutli and Mack, 1991); in Germany, as predator of tomato pests (Probst *et al.*, 1999); in Spain, as a predator of maize aphid (Asin and Pons, 1998). On the other hand, Johri and Reeta (1996), in India, observed that *L. riparia* form *bengalensis* (Dohrn) was omnivorous and the adults damaged 10.4-21% of stored potato tubers. In, Egypt, *L. riparia* was recorded affecting the changes in the population densities of *Thrips tabaci* Lind., *Empoasca* sp. and *Compylomma* spp. in corn fields (Ahmed and Darwish, 1991); *Bemisia tabaci* Gennadius in cotton fields (Darwish and Farghal, 1990); *Phthorimaea operculella* Zeller in tomato and potato fields (Abbas *et al.*, 1993) and *Aphis craccivora* Koch. in faba bean fields (El-Defrawi *et al.*, 2000). However, scanty attention has been paid to use the earwig as a biological control agent for the RPW immature stages. Accordingly, we aimed to examine the predacious efficiency of *L. riparia* for controlling the red palm weevil immature stages.

## **MATERIALS AND METHODS**

### **1. Rearing technique of red palm weevil, *Rhynchophorus ferrugineus*:**

Adults of red palm weevil, *Rhynchophorus ferrugineus* (Olivier), were collected from the fields which cultivated with date palm trees located at El-Katta, Giza Governorate. Plastic boxes were prepared (40 x20x15 cm) covered with muslin cloth for aeration, provided with small pieces of sugarcane stem. Each box contained 20♀+10♂. Newly hatched larvae were collected daily by pieces of sugarcane introduced to larvae for feeding. The last larval instar prior to pupation was placed in plastic boxes as mentioned above with palm pests. In addition, of large piece of sugarcane was used to help in forming the cocoons. Cocoons were collected, kept in bigger plastic boxes and supplied with wet filter paper. The obtained cocoons were daily inspected and emerged adults were collected and sexed soon after emergence. Paired adults placed in separate cups and provided with slices of sugarcane stem as ovipositional sites.

**2. Rearing of the earwig, *Labidura riparia*:**

Newly collected earwigs were reared on *Spodoptera littoralis* Boisduval larvae. Insects were kept in cylindrical plastic containers measuring 12cm. in diameter and 7cm. high containing 5cm. layer of moisturized sand. The containers were covered with halves of Petri dishes or with pieces of cloth fitted in position by rubber bands. For studying the biological processes, the individuals of earwigs were kept between two glass-plates that slipped in a woodier frame (20x20cm.), leaving a space of 1.5cm. between them. This space was filled with moistened sand to a level of 15cm. of the erected side, leaving the rest of the space for the free movement of the insects. Biological processes, such as mating, oviposition, maternal care, etc. were easily followed through the glass plates. Nymphs were reared individually in 9 cm. Petri dishes; a moistened filter paper was placed on the bottom of the dish and a piece of polyethylene under the lid. Number of laid eggs per female was determined by confining couples of recently emerged adults in deep Petri (EL-Hussini, 1969). All earwig individuals were reared and incubated under laboratory conditions at 25±1° C and 70±5 % R.H. where daily inspection and renewal of preys was carried out.

**3. Data analysis:**

Means were tested for significance by the one way analysis of variance (ANOVA) using Costat statistics 6.0, release 6.303 software. When the ANOVA statistics were significant at (P ≤ 0.01), means were compared by the Duncan's multiple range test using Costat statistics 6.0.

**RESULTS AND DISCUSSION**

Data presented in Table (1) showed the predative efficacy of nymphal stages of *L. riparia*. The 6<sup>th</sup> nymphal stage was the most predacious stage. The mean number of consumed immature stages of RPW was 349.33± 34.90, 351.33± 34.54, 340.67± 33.34, and 363.67± 36.67, respectively. Coming in the second place was the 5<sup>th</sup> nymphal instar, followed by 4<sup>th</sup> and 3<sup>rd</sup> nymphal instar. The 1<sup>st</sup> and 2<sup>nd</sup> nymphal instars of *L. riparia* were the least consuming instars. Results also showed that the 1<sup>st</sup> and 3<sup>rd</sup> larval instars of the RPW were the most preferable for consuming by the earwig 1<sup>st</sup>, 2<sup>nd</sup>, and 6<sup>th</sup> nymphal instars. However, the 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, and 5<sup>th</sup> nymphal instar preferred the RPW eggs for consumption (Table 1).

**Table (1): Mean numbers of red palm weevil immature stages consumed by nymphal stage of *Labidura riparia* under laboratory conditions**

Nymphal instars	Consumed red palm weevil stages (mean ± S. E.)			
	Eggs	1 <sup>st</sup> larval instar	2 <sup>nd</sup> larval instar	3 <sup>rd</sup> larval instar
1 <sup>st</sup> instar	93.67±18.45 <sup>a</sup>	96.00±15.7 <sup>a</sup>	85.33±12.33 <sup>b</sup>	80.33±13.30 <sup>c</sup>
2 <sup>nd</sup> instar	167.67±20.90 <sup>a</sup>	155.00±16.87 <sup>b</sup>	155.33±14.78 <sup>b</sup>	155.33±14.78 <sup>b</sup>
3 <sup>rd</sup> instar	219.67±25.8 <sup>a</sup>	214.33±22.88 <sup>a</sup>	174.00±16.67 <sup>b</sup>	171.00±16.01 <sup>b</sup>
4 <sup>th</sup> instar	253.33±33.3 <sup>a</sup>	149.00±10.33 <sup>b</sup>	152.00±15.78 <sup>b</sup>	115.33±11.25 <sup>c</sup>
5 <sup>th</sup> instar	332.67±33.33 <sup>a</sup>	331.00±32.03 <sup>a</sup>	323.67±30.67 <sup>a</sup>	273.33±25.33 <sup>b</sup>
6 <sup>th</sup> instar	349.33±34.90 <sup>a</sup>	351.33±34.54 <sup>a</sup>	340.67±33.34 <sup>a</sup>	363.67±36.67 <sup>a</sup>

**Table (2): Mean feeding duration of different nymphal stages of *Labidura riparia* on immature stage of red palm weevil under laboratory conditions**

Nymphal instars	Duration (Days) (mean $\pm$ S. E.)			
	Eggs	1 <sup>st</sup> larval instar	2 <sup>nd</sup> larval instar	3 <sup>rd</sup> larval instar
1 <sup>st</sup> instar	15.43 $\pm$ 1.48 <sup>a</sup>	15.89 $\pm$ 1.91 <sup>a</sup>	12.19 $\pm$ 2.09 <sup>b</sup>	11.47 $\pm$ 1.93 <sup>b</sup>
2 <sup>nd</sup> instar	15.14 $\pm$ 0.74 <sup>a</sup>	14.03 $\pm$ 0.72 <sup>a</sup>	12.94 $\pm$ 1.27 <sup>b</sup>	11.14 $\pm$ 0.78 <sup>c</sup>
3 <sup>rd</sup> instar	15.23 $\pm$ 0.98 <sup>a</sup>	14.30 $\pm$ 1.71 <sup>a</sup>	12.11 $\pm$ 2.04 <sup>b</sup>	11.08 $\pm$ 1.09 <sup>c</sup>
4 <sup>th</sup> instar	16.55 $\pm$ 1.48 <sup>a</sup>	15.03 $\pm$ 0.64 <sup>a</sup>	13.33 $\pm$ 2.01 <sup>b</sup>	12.22 $\pm$ 0.99 <sup>b</sup>
5 <sup>th</sup> instar	22.33 $\pm$ 1.98 <sup>a</sup>	17.89 $\pm$ 2.01 <sup>a</sup>	14.30 $\pm$ 1.22 <sup>b</sup>	13.00 $\pm$ 0.4 <sup>b</sup>
6 <sup>th</sup> instar	25.49 $\pm$ 1.94 <sup>b</sup>	30.33 $\pm$ 0.77 <sup>a</sup>	28.89 $\pm$ 0.99 <sup>a</sup>	31.13 $\pm$ 2.06 <sup>a</sup>

Data in Table (2) showed the mean duration of different nymphal stages of *L. riparia* fed on immature stages of red palm weevil. The 6<sup>th</sup> instar had the longest feeding duration upon all immature stages of RPW. The earwig nymphs consumed RPW eggs for 25.49  $\pm$  1.94 days, 1<sup>st</sup> instar larvae for 30.33  $\pm$  0.77, 2<sup>nd</sup> instar larvae for 28.89  $\pm$  0.99, and 31.13  $\pm$  2.06 days on the 3<sup>rd</sup> instar larvae. followed by 5<sup>th</sup> nymphal instar consumed (22.33 $\pm$ 1.98<sup>a</sup> eggs, 17.89 $\pm$ 2.01<sup>a</sup> 1<sup>st</sup> larval instar, 14.30 $\pm$ 1.22<sup>b</sup> 2<sup>nd</sup> larval instar and 13.00 $\pm$ 0.4<sup>b</sup> 3<sup>rd</sup> larval instar, 4<sup>th</sup> nymphal instar consumed (16.55 $\pm$ 1.48<sup>a</sup> eggs, 15.03 $\pm$ 0.64<sup>a</sup> 1<sup>st</sup> larval instar, 13.33 $\pm$ 2.01<sup>b</sup> 2<sup>nd</sup> larval instar and 12.22 $\pm$ 0.99<sup>b</sup> 3<sup>rd</sup> larval instar ), 3<sup>rd</sup> nymphal instar 15.23 $\pm$ 0.98<sup>a</sup> eggs, 14.30 $\pm$ 1.71<sup>a</sup> 1<sup>st</sup> larval instar, 12.11 $\pm$ 2.04<sup>b</sup> 2<sup>nd</sup> larval instar and 11.08 $\pm$ 1.09<sup>c</sup> 3<sup>rd</sup> larval instar, 2<sup>nd</sup> nymphal instar 15.14 $\pm$ 0.74<sup>a</sup> eggs , 14.03 $\pm$ 0.72<sup>a</sup> 1<sup>st</sup> larval instar, 12.94 $\pm$ 1.27<sup>b</sup> 2<sup>nd</sup> larval instar and 11.14  $\pm$ 0.78<sup>c</sup> 3<sup>rd</sup> larval instar and the 1<sup>st</sup> nymphal instar (15.43 $\pm$ 1.48<sup>a</sup> eggs, 15.89 $\pm$ 1.91<sup>a</sup> 1<sup>st</sup> larval instar 12.19 $\pm$ 2.09<sup>b</sup> 2<sup>nd</sup> larval instar and 11.47 $\pm$  1.93<sup>b</sup> 3<sup>rd</sup> larval instar.

Generally, it was observed that the 6<sup>th</sup> nymphal instar *Labidura riparia* of can survive for 31.13 days when fed on the 3<sup>rd</sup> larval instar of RPW red palm weevil under laboratory conditions. 25 °C $\pm$ 2 & Rh 70%  $\pm$ 5% Results were consistent with Kharboutli and Mack (1993) who found that *L. riparia* consumed more larvae of *Spodoptera frugiperda* (Smith) at 30 °C. Furthermore, the obtained results agreed with Bassal *et al.* (2001) who found that the rate of consumption of earwig nymphs increased by nymphal instar. In addition, Abd-Elgayed and Owayss (2007) in full agreement with our results. So, in conclusion, the earwig, *Labidura riparia*, as a predator, can be considered as a promising natural enemy to minimize the red palm weevil population density.

**Table (3): Nymphal duration of *Labidura riparia* fed on different immature stages of RPW.**

Nymphal instars	Eggs	1st larval instar	2nd larval instar	3rd larval instar
1 <sup>st</sup> instar	6 $\pm$ 1	6.5 $\pm$ 0.5	4.5 $\pm$ 0.5	4 $\pm$ 1
2 <sup>nd</sup> instar	11 $\pm$ 2	10.5 $\pm$ 1.5	10.5 $\pm$ 2.5	11.5 $\pm$ 2.5
3 <sup>rd</sup> instar	11 $\pm$ 2	13.5 $\pm$ 0.5	10.5 $\pm$ 0.5	14.5 $\pm$ 0.5
4 <sup>th</sup> instar	9 $\pm$ 0.5	12.5 $\pm$ 0.5	13 $\pm$ 1	12.5 $\pm$ 2.5
5 <sup>th</sup> instar	11.5 $\pm$ 0.5	13.5 $\pm$ 1.5	11 $\pm$ 1	13 $\pm$ 2
6 <sup>th</sup> instar	14.5 $\pm$ 0.5	14 $\pm$ 1	13.5 $\pm$ 0.5	10.5 $\pm$ 0.5

Nymphal duration of *Labidura riparia* fed on eggs was (70.5days) following by 1<sup>st</sup> instar larvae of RPW (66 days), the 2<sup>nd</sup> instar larvae (63days) and 3<sup>rd</sup> instar larvae (63 days).

## REFERENCES

- Abbas, M. S.T.; Abou-Zeid, N. A.; and Megahed, M. M. (1993). On the natural enemies of the potato tuber moth, *Phthorimaea operculella* in Egypt. Egyptian J. Agric. Res., 71 (4): 943-950.
- Abd-Elgayed, A. A. and Owayss, A. A. (2007). *Labidura riparia* Pallas (Dermaptera: Labiduridae) inhabiting honey bee colonies. Annals. Agric. Sci., Ain Shams Univ., Vol. 52(1): 243 – 251.
- Abraham, V. A.; Al-Shuaibi, M. A.; Faleir, J. R.; Abouzhairah, R. A.; and Vidyasagar, P. S. P.V. (1998). An integrated management approach for red palm weevil, *Rhynchophorus ferrugineus* Olivier, a key pest of date palm in the Middle East. Sultan Qabus Univ., J. Sci. Res. Agric. Sci., 3: 77-84.
- Ahmed, S.A. and Darwish, Y. A. (1991). Influence of weed control on sucking insect pests and natural enemies in a corn field in Assiut. Assiut J. Agric. Sci., 22 (2): 3-13.
- Asin, L. and Pons, X. (1998). Aphids predators in maize fields. Bull. OILB/SROP, 21 (8): 163-170.
- Bassal, T. T. M.; El-Naggar, M. E.; Fahmy, N. M.; Dorrah, M. A. A.; Sallam, M. H.; Salama, M. S. (2001). Carnivory, rate of digestion, and prey consumption by *Labidura riparia* (Dermaptera: Labiduridae). EFFLATOUNIA, 1: 13-19.
- Darwish, Y. A. and Farghal, A. I. (1990). Evaluation of certain aspesticides activity against the cotton whitefly, *Bemisia tabaci* and associated natural enemies on cotton plants under field conditions in Assiut. Assiut J. Agric. Sci., 21 (5): 331-339.
- Domiciano, N. L. and Herzog, D. C. (1990). Population dynamics of pests and some of their predators in soybean field under influence of herbicidal applications of Toxaphene. Resquisia Agropecuaria, Brasileria, 25 (2): 253-273.
- Dugard, R. (2006). Ed. Orange County Vector Control District. Pest Control Bulletin No. 13. 13001 Garden Grove Boulevard, CA 92843. [www.ocvcd.org](http://www.ocvcd.org).
- El-Defrawi, G.M.; Emam, A. K.; Marzouk, I. A.; and Rizkala, L. (2000). Population dynamics and seasonal distribution of *Aphis craccivora* K. and associated natural enemies in relation to virus disease incidence in faba bean field. Egyptian J. Agric. Res., 78 (2): 627-641.
- El-Hussieni, M. M. (1969). Studies on the Earwig, *Labidura riparia* Pallas. M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt. Pp. 206.

- Girgis, G. N.; Batt, A. M.; Okil, A. M.; Haggag, S. M.; and Abdel Azim, M. M. (2002). Evaluation of trunk injection methods for the control of red palm weevil, *Rhynchophorus ferrugineus* (Oliver) in date palm trees in Egypt. 2<sup>nd</sup> International Conference, Plant Protection Institute, Cairo, Egypt, 21-24 December 2002; 1: 709-711.
- Guimaraes, J.H.; Tocci, E. C.; and Gomes, J. P. C. (1992). Dermaptera (Insecta) associated with commercial poultry ranches in Sao Paulo State and their importance as biological control agents of poultry pests. Rev. Brasil. Entomol.; 36(3): 527-534.
- Johri, P. K. and Reeta, J. (1996). Biology and extent of damage of the Indian predaceous earwig *Labidura riparia* Pallas form bengalensis (Dohrn) (Dermaptera: Labiduridae) as an occasional pest of potato tuber in Uttar Pradesh. Bioved, 7 (1): 11-21.
- Kharboutli, M. S. and Mack, T. P. (1991). Relative and seasonal abundance of predaceous arthropods in Alabama peanut fields as indexed by Pitfall traps. J. Econ. Entomol.; 84(3):1015-1023.
- Kharboutli, M. S. and Mack, T. P. (1993). Effect of temperature, humidity and prey density on feeding rate of the striped earwig (Dermaptera: Labiduridae). Environ. Entomol.; 22(5): 1134-1139.
- Lee, J. H.; Johnson, S. J.; and Wright, V. L. (1990). Quantitative survivorship analysis of the velvetbean caterpillar (Lepidoptera: Noctuidae) pupae in soybean fields in Louisiana, USA. Environ. Entomol.; 19(4): 978-986.
- Murphy, S. and Briscoe, B. (1999). The red palm weevil as an alien invasive: biology and prospects for biological control as component of IPM. Biocontrol News and Information, 20(1): 35-46.
- Probst, K.; Pulschen, L.; Sauerborn, J.; and Zebitz, C. P. W. (1999). Influence of seasonal pesticide use regimes on the entomofauna of tomato on highland in Ecuador. Maejo. Integrado de Plagas, 45: 53-62.
- Wattanapongsiri, A. (1966). A revision of the genera *Rhynchophorus* and *Dynamis* (Coleoptera: Curculionidae). Bangkok, Thailand, Dept. Agric. Sci. Bull., 328 p.

### الكفاءة الافتراسية لإبرة العجوز الكبيرة *Labidura riparia* المرابة على الأطوار الغير الكاملة لسوسة النخيل الحمراء

عادل حسن عبد السلام<sup>١</sup> وعبد المنعم عبد الودود البنأ<sup>٢</sup> و ايمان السيد حسين الرهيوى<sup>٣</sup>  
جامعة المنصورة - كلية الزراعة - قسم الحشرات الاقتصادية<sup>١</sup>  
معمل بحوث تطوير النخيل- مركز البحوث الزراعية- الدقى- جيزة- مصر<sup>٢</sup>  
معهد بحوث وقاية النباتات - الدقى -جيزة- مصر<sup>٣</sup>

تم إجراء الدراسة الحالية معملياً لتقييم كفاءة النشاط الافتراسى لإبرة العجوز ضد الأطوار الغير كاملة لسوسة النخيل الحمراء. أظهرت النتائج المتحصل عليها أن الحوريات فى العمر السادس لإبرة العجوز كان الأكثر افتراساً للأطوار الغير كاملة. وكان أقلهم افتراساً هو العمر الحورى الأول لإبرة العجوز وأقل معدل افتراس. أوضحت النتائج أن معدل الافتراس لأطوار الغير كاملة لسوسة النخيل تزيد بزيادة العمر الحورى لإبرة العجوز. كما أوضحت النتائج أيضاً أن العمر الحورى السادس استغرق أطول مدة افتراس على الأطوار الغير كاملة لسوسة النخيل. تعتبر ابرة العجوز عدو طبيعى واعد فى تقليل أعداد سوسة النخيل الحمراء.