

14- Vasectomy failure rate is quoted as approximately 1 in 2000 (0.05%) after clearance has been given. By how many months post procedure should the vasectomy be considered a failure if motile sperm are still observed in a fresh semen sample?

- A. 3 months
- B. 4 months
- C. 5 months
- D. 6 months
- E. 7 months

15- Ulipristal acetate (UPA) may itself reduce the efficacy of other hormonal contraception. When emergency contraception is administered because of a missed combined oral contraceptive pill (CHC) the CHC should be continued. For how many days should additional contraception be advised?

- A. 0 days
- B. 7 days
- C. 14 days
- D. 21 days
- E. 28 days

16- As a post-coital contraception, the primary mechanism of action of Ulipristal acetate is inhibition of ovulation. What is the conception rate when Ulipristal acetate is taken within 120 hours of unprotected sexual intercourse?

- A. 1.3%
- B. 2.3%
- C. 3.3%
- D. 4.3%
- E. 5.3%

17- A 28-year-old patient is referred from the A&E department with lower abdominal pain and vaginal discharge. On examination she has a temperature of 37.3°C, a pulse of 86 bpm and a BP of 120/83 mm Hg. Her abdomen is slightly tender in the LIF region. On vaginal examination there is a mild corresponding adnexal tenderness, the IUCD threads are seen and a swab of mucopurulent discharge is taken. She mentions that she has a new sexual partner and has not used barrier contraception. What is the most appropriate next step in her management?

Options

- A. Intravenous PID-targeted antibiotics + IUCD removal immediately.
- B. Oral PID-targeted antibiotics + IUCD removal immediately.
- C. Oral PID-targeted antibiotics and, if no improvement in symptoms, remove IUCD in 24 hours.
- D. Oral PID-targeted antibiotics and, if no improvement in symptoms, remove IUCD in 48 hours.
- E. Oral PID-targeted antibiotics and, if no improvement in symptoms, remove IUCD in 72 hours.