

31. A lens coloboma:

- a. Is usually associated with previous lens trauma
- b. Is typically located superiorly
- c. Is typically associated with normal zonular attachments
- d. Is often associated with cortical lens opacification

32. Three years old child with esotropia 30 prism diopters, cycloplegic refraction right eye +2.00 DS, left eye +0.50 DS what do you do?

- a) Glasses full correction
- b) Surgery
- c) Atropine left eye
- d) Partial time closure of the left eye

33. Cyclosporine

- a) Cytotoxic antimetabolite
- b) Cytostatic anti-inflammatory
- c) Immune modulator of interleukin 2
- d) Cytotoxic alkylating agent

34. What is the most common cause of bleb failure?

- a) Bleb encapsulation (Tenon cyst formation)
- b) Episcleral fibrosis
- c) Late bleb leak
- d) Closure of the internal sclerostomy

35. Which of the following systemic diseases is associated with central pigmentary macular changes?

- a) homocystinuria
- b) Ehlers-Danlos syndrome
- c) Marfan syndrome
- d) myotonic dystrophy

36. In patients with upper eye lid ptosis, which of the following is the most important measure in determining the type of surgery?

- a) Margin reflex distance
- b) Eye lid excursion
- c) Response to phenylephrine
- d) Pretarsal show

37. Least in management of postoperative endophthalmitis:

- a) Blood culture
- b) Systemic antibiotics
- c) Intravitreal antibiotics
- d) Intravitreal corticosteroids