

Effect of Antennae on the Honey Bee Queens Behavior Inside and Outside of their Hives

Sawsan M. Abdel-Megeed

Plant Protection Department, Faculty of Agriculture, Ain Shams University, Cairo, Egypt

Corresponding authors, E. mail: sawsangida@yahoo.com



ABSTRACT

The present study was carried out during season 2014 using honey bee colonies headed with open mated local Italian queens, *Apis mellifera lugustica*, at the apiary belonging to Faculty of Agriculture, Ain Shams University, to explain the role of antennae of queens in controlling or managing their behavior. The experiment was done by using 12 virgin or mated queens. Each of them was divided equally into four treatment i.e queens with only one flagellum, queens without the last segments of both flagella, queens without both flagella and control. The results showed that the virgin queens with one (single) flagellum could be mated and laid eggs regularly but the virgin queens without flagellum did not leave their hive. On the other hand, when the last segment of both flagella was cut in virgin queens, they could not return to their hive and may be lost. The mated queens with one (single) flagellum and without the last segment (No 10) of flagellum could lay their eggs regularly in combs as the control but mated queens without flagellum stopped laying eggs for one week then began to lay eggs unregularly in the comb, lay two eggs in the same cell and put the drone's eggs in small cells (25 cells/ inch²). The number of sealed brood cells in the first and second treatments was significantly as the same as control (mated queen with single flagellum and mated queens without the last segment of flagellum). On contrast, the mated queens without both flagella had significantly less numbers of brood cells than other treatments.

Keywords: Honey bee queen– *Apis mellifera* – Flagellum – Sense organs – Behavior .

INTRODUCTION

The antenna of honey bees carries a lot of sense organs (Abdelmegeed & Sawires, 2015; Al-Ghamdi, 2006; Naik *et al*, 1995) such as mechanoreceptors (trichoid and campaniform) and chemical sense organs such as placoid (Getz and Akers, 1994) and basiconicum and different diameter of pores such as sensilla coeloconica and sensilla ampullacea (Stort and Malaspina, 1997) which affected on behavior of honey bee workers. (Jung *et al.*, (2014); Rogers *et al*, (2013); Muenz *et al*, (2012); Rigosi *et al*, (2011) studied the effect of sense organs on the behavior of bees (i.e. hive land marker, hygienic (Gramacho *et al*, 2003), pollen & nectar collecting). (Tsuruda and Page (2009); Sanchez *et al* (2005); Haupt (2004) also studied the defense of bee about their hive and behavior of queens (i.e. mating and method of laying the eggs regularly. So that, the antenna is play an important role in behavior of honey bees.

Therefore, the present study aimed to explain the role of antennae in affecting behavior of honey bee queens inside and outside their hives.

MATERIALS AND METHODS

The experiments of the present work were conducted at the apiary belonging to Faculty of Agriculture, Ain Shams University. The used bee colonies were headed with open mated local Italian queens, *Apis mellifera lugustica*

The queens were divided into two parts (virgin queens or mated queens) each part contained 12 queens divided to four groups each one consisted of three queens:-

The first group: contained queens with one flagellum (virgin or mated) by removing or cutting the other one from the pedicel part.

The second group: contained queens (virgin or mated) without the tip of flagellum by cutting segments number 10.

The third group: contained queens without both flagella by removing both of them from the pedicel part.

The fourth group: contained normal queens with pair of flagella (control).

All the treated queens (virgin or mated) were introduced to colonies each colony consists of three combs (two closed brood combs + one honey and pollen comb) in spring season.

Number of sealed brood cells

Number of sealed brood cells in colonies with queens at all treatments was calculated every 13 days according to :

$$\text{Area of sealed brood} = \pi \times r1 \times r2$$

$$\pi = 22/7$$

$$\text{Number of sealed brood cells} = \text{area of close brood} \times 3.875$$
$$3.875 = \text{number of cells}/\text{cm}^2$$

Statistical analysis

Chi Square (Fisher's Exact Test) and Tukey test (ANOVA) were used by using SAS program (2005).

RESULTS AND DISCUSSION

• The virgin queens:

From Table 1, it can be concluded that all virgin queens with single flagellum went outside their hives for mating and returned to their hives. They laid the eggs regularly in the combs, one egg inside each cell. The worker's eggs were laid inside small cells (25 cells/ inch²) and drone's eggs were inside large cells (16 cells / inch²). All virgin queens without last segment of flagella (segment number 10) went outside their hives; two queens did not return to their hives and lost but one queen could return to its hive but still virgin. All virgin queens without pair of flagella stayed inside the hives and were still virgin. In the control (normal virgin queens), all the virgin queens went out their hives, returned and put the eggs regularly.

Table 1. Behavior of treated virgin queens in mating

treatments	Behavior	Exit & mated	Exit & losted	Not mated
1		3	-	-
2		-	2	1
3		-	-	3
4		3	-	-
		Fisher's Exact Test:-	2.165E-04	
		Pr <= P	0.0026	
		Contingency Coefficient:-	0.7746	

- *(1) The virgin queens with single flagellum.
- *(2) Queens without the segment number 10 of the two flagella.
- *(3) Queens without pair of flagella.
- *(4) Normal queens (control).

• **The mated queens:**

Data arranged in Table 2, showed that, the mated queens with single flagellum laid their eggs regularly in the combs, put one egg inside one cell and acted normal (the worker's eggs put in small cells (25 cells/ inch²) and drone's eggs in large cells (16 cells/inch²). The mated queens without last segment of flagellum (segments number 10) acted normal as the previous treatment and control while, the mated queens without pair of flagella stopped putting eggs for one week and acts in abnormal behavior, or odd manner. The queens

moved inside the hive quickly under stress and looked inside the cell and again looked inside the other cell but did not put the eggs inside the cells (Fig. 1). Week after, the queens started to put the eggs not regularly in the comb. They put more than one egg inside one cell and the drone's eggs in small cells (25 cells/inch²) (Fig. 2). The mated untreated queens (control) acted normal in their behavior, all the queens put the eggs regularly in the comb, put one egg inside one cell, put the worker's eggs in small cells (25 cells/ inch²) and drone's eggs in large cells (16 cells/inch²).

Table 2. Behavior of treated mated queens in laying their eggs.

treatment	Behavior	eggs laid in the same day of cutting	Eggs laid week after cutting
1		3	0
2		3	0
3		0	3
4		3	0
		Fisher's Exact Test:-	0.0045
		Pr <= P	0.0182
		Contingency Coefficient :-	0.7071

- *(1) Queens with single flagellum.
- *(2) Queens without the last segment of flagellum (segment number 10).
- *(3) Queens without pair of flagella.
- *(4) Normal queens (control).




Fig 1. Parts of video showed the odd behavior of mated queens after cutting the pair of flagella in laying eggs in their hives.

In the present work, role of flagellum is very important where they affected on behavior of queens honey bee inside their hives such as laying eggs regularly and outside their hives such as mating and land marker their hives. Jung *et al* (2014) found that the olfactory organs are important sensory modalities on individual discrimination, perception and efficient orientation to get nectar and pollen in honey bees. Rogers *et al* (2013) showed that the honeybees preferred to use their right flagellum in social interactions. Gil & Marco (2010) studied the tactile hair on flagellum, enables bees to estimate the direction relative to gravity and the length of the waggle phase (waggle dance).

Number of sealed brood cells

The data given in Table 3 clearly showed that mean number of brood cells was affected by present or absent of flagellum in different treated queens.

In the beginning of experiments, the number of sealed brood cells did not significantly differ in all treatments and control. In second date (5/10/2014) the number of sealed brood cells was significantly differ between the first two treatments and control from one side and queens without both flagella from other sides which have no sealed brood cells. The number of the sealed brood cells at the second date was 7002.0, 6942.3 and 6942.3 in the control, queens with one flagellum and queens without the last segment of flagellum, respectively in comparison with queens without pair of flagella (0.00). In the third and fourth dates the number of brood cells was significantly higher in the queens with pair of flagella(control), with single flagellum and without the last segment of flagellum than queens without pair of flagella.

Table 3. Number of sealed brood cells/hive produced from treated mated queens every 13 days.

treatments	No. sealed brood cells/colony				
	Dates	22/9	5/10	18/10	31/10
1		7276.7 ±91.3175	6942.3 ^a ±73.7126	5785.3 ^a ±87.4274	5389.0 ^a ±56.8741
2		7459.7 ±130.2314	6729.0 ^a ±96.8538	5548.3 ^a ±132.2229	5223.7 ^a ±57.7888
3		7539.0 ±79.4858	0.0 ^b ±0.0000	1828.0 ^b ±104.5498	1833.3 ^b ±63.6884
CONT		7612.7 ±149.3906	7002.0 ^a ±131.4762	5876.7 ^a ±80.3008	5481.3 ^a ±128.3883
F.Value.		3.09	2962.12**	722.5**	928.59**
L.S.D.			286.87	330.11	262.86

:*means with the same superscript in the same column were not significantly different) p(. . ° >

1. Queens with single flagellum.
2. Queens without last segment of flagellum.
3. Queens without pair of flagella.



Fig 2. Sealed brood cells produced from mated queens without flagellum at both the third and fourth dates.

REFERENCES

- Abdelmegeed S. M. and S. G. Sawires (2015). Comparative studies on the antennal sense organs of queen and worker honey bees, *Apis mellifera* L. Egypt. Acad. J. Biolog. Sci., 8(1): 65-75.
- Al-Ghamdi, A. A. (2006). Scanning electron microscopic studies on antennal sensilla organs of adult honey bee workers in genus *Apis* (Hymenoptera: Apidae). Bull. Ent. Soc. In Egypt, 83: 1-11.
- Getz, W. M. and R.P. Akers (1994). Honeybee olfactory sensilla behave as integrated processing units. Behav. and Neural Biology, 61(2):191-195.
- Gramacho, K.P.; L.S. Goncalves; A.C. Stort and A.B. Noronha (2003). Is the number of antennal plate organs (sensilla placodea) greater in hygienic than in non-hygienic Africanized honey bees?. Genetics and Molecular Res., 2(3):309-316.
- Haupt, S. S. (2004). Antennal sucrose perception in the honey bee (*Apis mellifera* L.): behaviour and electrophysiology. J. Comparative Physiology. A, Sensory, Neural, and Behavioral Physiology, 190(9):735-745.
- Jung J.W; K. W. Park; H. W. Oh and H. W. Kwon (2014). Structural and functional differences in the antennal olfactory system of worker honey bees of *Apis mellifera* and *Apis cerana*. J. Asia-Pacific Ento., 17(3):639-646.
- Naik, D.G.; A.S. Bhongle; A.H. Kapadi; M.C. Suryanarayana; S.S. Chawda and O.R. Chaudhary (1995). Antennal sensilla of adult worker *Apis cerana indica*. J. Apic. Res., 34(4):205-208.
- Muenz, T. S.; A. Maisonnasse; E. Plettner; Y. Conte and W. Rossler (2012). Sensory reception of the primer pheromone ethyl oleate. Naturwissenschaften, 99(5):421-425.
- Rigosi, E.; E. Frasnelli; C. Vinegoni; R. Antolini; G. Anfora; G. Vallortigara and A. Haase (2011). Searching for anatomical correlates of olfactory lateralization in the honeybee antennal lobes: a morphological and behavioural study. Behav. Brain. Res., 221(1):290-294.
- Rogers, L.J.; E. Rigosi; E. Frasnelli and G. Vallortigara (2013). A right antenna for social behaviour in honey bees. Scientific Reports, <http://dx.doi.org/10.1038/srep02045>.
- Sanchez, M. G. de B.; M. Giurfa; T.R. de P. Mota and M. Gauthier (2005). Electrophysiological and behavioural characterization of gustatory responses to antennal 'bitter' taste in honeybees. Euro. J. Neuroscience; 22(12):3161-3170.
- Stort, A. C. and O. Malaspina (1997). Differences in the number of sensilla coeloconica and sensilla ampullacea of segment 10 of the antennae of Africanized and Caucasian bees and of their F1 hybrids. Brazilian. J. Genetics. 20(2):189-192.
- Tsuruda, J. M. and R.E. Page(2009). The effects of foraging role and genotype on light and sucrose responsiveness in honey bees (*Apis mellifera* L.). Behav. Brain Res., 205(1):132-137.

تأثير قرون الاستشعار على سلوك ملكات نحل العسل داخل وخارج الخلية

سوسن محمد عبد المجيد

قسم وقاية النبات - كلية الزراعة جامعة عين شمس - القاهرة - مصر

تم دراسة تأثير قرون الاستشعار للملكات في منحل كلية الزراعة جامعة عين شمس خلال موسم الربيع 2014 في طوائف ذات ملكات هجين ايطالى تم تلقيحها طبيعيا. التجارب أجريت باستخدام ١٢ ملكة عذراء أو ملقحة. كل واحدة منهم قسمت بالتساوى الى أربع معاملات (ملكات ذات قرن استشعار واحد و ملكات بدون العقلة الأخيرة من السوط و ملكات بدون قرون استشعار وكنترول). أظهرت النتائج أن الملكات العذارى التي لها قرن استشعار واحد قادرة على التزاوج ووضع البيض بانتظام داخل خليتها بينما الملكات العذارى التي تم قطع العقلة العاشرة (الطرفية) من قرن الاستشعار خرجت للتزاوج ولم تعود مما يدل على انها تاهت عن خليتها بينما الملكات العذارى التي تم قطع السوط بالكامل من زوج قرن الاستشعار لم تستطع الخروج من الخلية وظلت بداخلها وذلك بالمقارنة مع الملكات العذارى التي بها زوج من قرن الاستشعار كامل (كونترول) خرجت وتلقحت وعادت الى خليتها ووضع البيض بانتظام. كما أظهرت النتائج أن الملكات الملقحة التي لها قرن استشعار واحد والملكات الملقحة التي تم قطع العقلة العاشرة للسوط والملكات التي بها زوج من قرن الاستشعار كامل (كونترول) قادرين على وضع البيض بانتظام داخل الخلية بالإضافة الى وضع بيضة واحدة داخل كل عين ووضع بيض الذكور في عيون سداسية واسعة وبيض الشغالات داخل عيون سداسية ضيقة. أما الملكات الملقحة التي تم قطع السوط بالكامل من قرني الاستشعار توقفت عن وضع البيض لمدة أسبوع واصبحت متوترة وتحرك حركة سريعة وتنتظر داخل العين السداسية وتخرج دون ان تضع بيض ثم تعاود النظر مرة أخرى داخل عين سداسية أخرى وهكذا دون ان تضع بيض وبعد أسبوع بدأت في وضع البيض ولكن بدون انتظام ومبعثر على القرص وتضع أكثر من بيضة داخل العين السداسية الواحدة بالإضافة الى انها تضع بيض الذكور في عيون سداسية ضيقة. تساوت أعداد الحضنة المقفولة في كل من المعاملة الأولى (ملكات ملقحة نو قرن استشعار واحد) والمعاملة الثانية (ملكات ملقحة بدون العقلة الطرفية لقرني الاستشعار) مع الكنترول بينما انخفضت اعداد الحضنة في المعاملة الثالثة (ملكات ملقحة بدون قرني الاستشعار) انخفاض معنوى واضح عن بقية المعاملات و الكنترول.