

EFFECTS OF AIR POLLUTION ON MORPHOLOGICAL AND PHYSIOLOGICAL FEATURES OF *Malva parviflora*

Abu Ziada, M.E.A.; S.A. Haroun; G.A. El-Sherbeny and A. A. Najm

Botany Department, Faculty of Science, Mansoura University, Egypt.



ABSTRACT

Increasing industrial activity intensifies the release of different types of harmful pollutants into the atmosphere. Air pollution with dust, smoke and chemical vapor is hazardous to humans as well as to plants. This study describes the effect of air pollution resulted from brick factories and road side pollution on some morphological and physiological characteristics of *Malva parviflora* growing naturally around these pollution sources. The morphological traits as stem length, branch number, root length, leaf length, leaf width and leaf area as well as some physiological characteristics as levels of chlorophyll a, chlorophyll b, carotenoids and total soluble sugars were investigated. The results showed that the morphological and physiological features of *Malva parviflora* were reduced in the polluted areas in comparison with regions away from these areas.

Keywords: Air pollution, morphological traits, physiological characteristics, *Malva parviflora*.

INTRODUCTION

Pollutants released from different industries adversely affect man's food supply. The presences of such toxic pollutants in the environment are responsible for altering the ecosystem. The air pollution due to traffics and brick factories plays a significant role in the environment imbalances and in production of air pollution hazardous effects (Stern, 1976; Thambavani and kumar 2011; Thambavani and kumar 2012).

Air pollution causes diverse effects on the physiological processes in plants that lead to reduction in plant growth (Schutzki and Cregg, 2007). Air pollution affects many metabolic processes in plants such as photosynthesis, mitochondrial respiration and stomatal clogging (Miller *et al.*, 1973).

Taylor and Davies (1990) reported that the ambient level of air pollution affects stomatal conductance, photosynthesis and root morphology of young beech. Out of the most recent studies indicated that dust accumulation causes severe damage in the photosynthetic apparatus is study by Santosh and Tripathi (2008).

Besides causing suppression of plant growth, pollution dust induces the changes in the physico-chemical properties of the soil rendering it (Parthasarthy *et al.*, 1975; Singh and Rao, 1978; Addo *et al.*, 2013). In comparison to gaseous air pollutants, only limited studies have been carried out on the effect of particulate air pollutants on plant including fluoride dust (McCune *et al.*, 1965), soot (Miller and Rich, 1967), lead particles, brick factories pollutants, roadside pollutants, cement dust (Darley, 1966; Pandey and Simba, 1988, 1990) and coal dust (Rao, 1971).

The aim of the study was to investigate the impacts of roadside and brick factories pollutants on morphological characteristics and physiological features of *Malva parviflora*.

MATERIALS AND METHODS

The *Malva parviflora* samples growing naturally around road sides and brick factories were collected at zero meter (polluted sites) and 120 meters (non-polluted sites) from the studied pollution sources.

Chlorophyll a, chlorophyll b and carotenoids were analyzed following the method of Lichtenthaler (1987). Total soluble sugar was determined by the method adopted by Thayumanavan and Sadasivam (1984). Leaf area was estimated as Length X Breadth X 0.794 (Carvalho *et al.*, 2011). The samples were investigated in triplicate.

RESULTS AND DISCUSSION

The morphological characteristics:

The obtained results showed significant reduction in the morphological characteristics in the polluted sites at roadside (Table 1) than the sites at 120 meters away from these sites (considered as control). Where the stem length was 28.13 ± 4.99 cm at the polluted sites and 41.25 ± 8.26 cm at 120 meters away from polluted sites; root length was 10.28 ± 1.37 cm at the polluted sites and 12.94 ± 1.70 cm at 120 meters. Also, number of branches, leaf length, leaf width and leaf area were recorded in polluted sites and at 120 meters as follow: 9.5 ± 3.16 and 16.88 ± 5.28 ; 4.68 ± 0.31 cm and 6.36 ± 0.63 cm respectively leaf width from 5.2 ± 0.46 and 7.33 ± 0.77 as well as 18.24 ± 2.11 cm² and 34.98 ± 5.04 cm² respectively.

Meanwhile, at the brick factories there were significant reduction in the morphological characteristics at the polluted sites as stem length from 42.50 ± 4.50 cm at the factory sites to 24.25 ± 4.03 cm at 120 meters away from factory sites, root length from 8.23 ± 1.67 cm at the factory sites to 6.30 ± 1.46 cm at 120 meters away from factory sites, number of branches from 19.00 ± 3.46 at the factory sites to 11.00 ± 3.16 at 120 meters away from factory sites, leaf length from 5.85 ± 0.57 cm at the factory sites to 4.08 ± 0.61 cm at 120 meters away from factory sites, leaf width from 4.40 ± 0.89 cm at the factory sites to 2.75 ± 0.37 cm at 120 meters away from factory sites and leaf area from 19.57 ± 5.39 cm² at the factory sites to 8.48 ± 2.22 cm² at 120 meters away from factory sites (Table 1). The Reduction increased with increasing the distance from the factories where the dust and pollutants released from their chimneys are at high elevation and so carried by air to affect plants away the factory itself.

Generally, the leaf area, leaf length, leaf width, number of branches, stem and root length were significantly reduced in polluted area than in areas away from pollution. Air pollutants can cause leaf injury, stomatal damage, reduced photosynthetic activity, disruption in membrane permeability and reduction in the growth and yield of the plants. Previous research reported

significant reduction in many leaf growth parameters in the polluted areas in comparison with the control ones (Jahan and Iqbal, 1992; Seyyednejad *et al.*, 2011). The obvious reduction in the leaf surface area causes less contact with air pollutants and improves plant resistance to pollution. Reduction in leaf area of many plant species growing in the vicinity of heavy pollutants was observed. Some hidden injury or physiological disturbance might be occurred in addition to the reduction in morphological and anatomical attributes (Seyyednejad *et al.*, 2009). Long-lasting impact of different pollutants including sulphur and nitrogen oxides and heavy metals causes a reduction in the leaf size and growth of aerial parts (Kozlov *et al.*, 1999; Tiwari *et al.*, 2006). These leaf characteristics can demonstrate the air pollution (Jahan and Zafar, 1992).

Table 1. The comparison of morphological characteristics of *Malva parviflora* in the study sites.

Pollution source	Roadside		Brick factories	
	Polluted sites	120m away	Polluted sites	120m away
Stem length (cm)	28.13 ± 4.99	41.25 ± 8.26	42.5 ± 8.54	24.25 ± 4.03
Branches number	9.5 ± 3.16	16.88 ± 5.28	19.00 ± 3.46	11.00 ± 3.16
Root length (cm)	10.28 ± 1.37	12.94 ± 1.70	8.23 ± 1.67	6.3 ± 1.46
Leaf length (cm)	4.68 ± 0.31	6.36 ± 0.63	5.85 ± 0.57	4.08 ± 0.62
Leaf width (cm)	5.2 ± 0.46	7.33 ± 0.77	4.4 ± 0.89	2.75 ± 0.37
Leaf area (cm ²)	18.24 ± 2.11	34.98 ± 5.04	19.58 ± 5.39	8.48 ± 2.22

Polluted sites = at zero meter; 120m away= 120 meters away from the pollution source. Values are the means of replicates ± standard deviation.

The biochemical characteristics:

It is clear from the results that the leaves collected from roadside polluted areas and at 120 meters from polluted areas showed significant decrease in chlorophyll a from 2.70 ± 0.85 mg/g fresh weight at 120 meters away from polluted sites to 1.16 ± 0.15 mg/g fresh weight at the polluted sites, chlorophyll b from 1.54 ± 0.58 mg/g fresh weight to 0.57 ± 0.11 mg/g fresh weight, carotenoids from 0.82 ± 0.48 mg/g fresh weight to 0.28 ± 0.13mg/g fresh weight and soluble sugar from 146.9 to 70.9 mg/g fresh weight. While the leaves collected from brick factories polluted area showed significant decrease in chlorophyll-a from 1.60 ± 1.10 mg/g fresh weigh at the factory sites to 1.17 ± 0.16 mg/g fresh weight at 120 meters from the factory sites, chlorophyll b from 0.86 ± 0.58 mg/g fresh weight at the factory sites to 0.65 ± 0.12 mg/g fresh weight at 120 meters away from the factory sites, carotenoids from 0.65±0.11 mg/g fresh weight at the factory sites to 0.29 ± 0.13 mg/g fresh weight at 120 meters from factory sites and soluble sugar from 128.88 at factory sites to 75.18 mg/g fresh weight at 120 meters away from factory sites (Table 2). The Reduction increased with increasing the distance from the factories where the dust and pollutants released from their

chimneys are at high elevation and so carried by air to affect plants away the factory itself.

Chlorophyll and carotenoids are the main core of energy production in green plants, and their amounts are significantly changed by environmental effects on plant metabolism (Shweta and Agrawal, 2006). The photosynthetic pigments are most likely to be damaged by air pollution. Chlorophyll pigments under stress might undergo several photochemical reactions such as oxidation, reduction, phaeophytinisation and reversible bleaching (Giri *et al*, 2015). The chloroplast damaged by incorporation of cement dust on leaf caused reduction in chlorophyll concentration in the plants which are near the industry (Lerman, 1975; Singh and Rao, 1978). Similar results were observed in maize (Pandey and Simba, 1989; Pandey *et al.*, 1999). Sulphur oxides and nitrogen oxides have detrimental effects that degrade chlorophyll to a photosynthetically inactive phaeophytin and Mg⁺⁺. Any alteration in chlorophyll concentration may change the morphological, physiological and biochemical behaviour of the plant. Air pollution-induced degradation in photosynthetic pigments was also observed (Bansal, 1988; Singh, 1990; Sandelius, 1995, Giri *et al*, 2013). Many researchers reported reduced carotenoids content under the effect of air pollution (Joshi *et al*, 2009; Tiwari *et al*, 2006; Giri *et al*, 2013).

In plant tissue, soluble sugars have osmoprotectant and cryoprotectant roles and their presence is important for plasma membrane integrity. Soluble sugars are important components in plant tissues; represent the source of energy in all organisms (Shvaleva *et al*, 2005; Moraga *et al*, 2006; Naya *et al*, 2007). Soluble sugars significantly reduced under air pollution due to the increase in respiration and decrease of CO₂ fixation because of chlorophyll deterioration. Pollutants like sulphur oxides and nitrogen oxides might cause more depletion of total soluble sugars in leaves of plants grown in polluted areas (Tripathi and Gutam, 2007; Giri *et al*, 2013).

Table 2. The comparison of biochemical characteristics of *Malva parviflora* in the study sites.

Pollution source distance	Roadside		Brick factories	
	Polluted sites	120m away	Polluted sites	120m away
Chlorophyll a mg/g FW	1.16 ± 0.15	3.70 ± 0.85	1.60 ± 1.10	1.17 ± 0.16
Chlorophyll b mg/g FW	0.57 ± 0.11	1.21 ± 0.58	0.86 ± 0.58	0.68 ± 0.13
Carotenoid mg/g FW	0.28 ± 0.13	0.82 ± 0.48	0.65±0.11	0.29 ± 0.13
Total soluble sugars mg/g DW	70.9	146.9	128.88	75.18

Polluted sites = at zero meter; 120m away= 120 meters away from the pollution source.
FW= Fresh weight, DW= dry weight.

CONCLUSION

The concentrations of chlorophyll a, chlorophyll b, carotenoids and total soluble sugar in *Malva parviflora* reduced in polluted regions. In addition, the growth traits of *Malva parviflora* were reduced in polluted regions compared with regions away from pollution.

REFERENCES

- Addo M. A., Darko E. O. and Gordon C. (2013). Contamination of soils and loss of productivity of Cowpea (*Vigna unguiculata* L.) caused by cement dust pollution. *International Journal of Research in Chemistry and Environment*, 3(1): 272-282.
- Bansal S. (1988). "Studies on the effect of certain atmospheric pollutants on fruit diseases of *Lycopersicon esculentum* Mill. Caused by *Alternaria alternate*", PhD thesis, Bhopal, Bhopal University.
- Darley E. F. (1966). Studies on the effect of cement kiln dust on vegetation. *Journal of the Air Pollution Control Association*, 16: 145-150.
- Carvalho, L.B.I; Souza, M.C.; Bianco, M.S. and Bianco, S. (2011). Estimate of the leaf area of aquatic weeds: *Pistia stratiotes*. *Planta daninha*, 29 (1): 65-68.
- Giri S., Shrivastava D., Deshmukh K. and Dubey P. (2015). Effect of Air Pollution on Chlorophyll Content of Leaves. *Current Agriculture Research Journal*, 1(2): 93-98.
- Jahan S. and Iqbal M. Z. (1992). Morphological and anatomical studies of leaves of different plants affected by motor vehicles exhaust. *Journal of Islamic Academy of Science*, 5:21-23.
- Joshi P.C. and Swami A. (2009). Air pollution induced changes in the photosynthetic pigments of selected plant species. *Journal of Environmental Biology*, 30: 295-298 .
- Kozlov M.V., Zvereva E.L. and Niemelä P. (1999). Effects of soil quality and air pollution on the rooting and survival of *Salix borealis* cuttings. *Boreal Environmental Research*, 4: 67–76.
- Lerman S. L. and Darley E. F. (1975). Particulates, In. *Responses of Plants to Air Pollution*. pp. 141. New York: Academic Press.
- Lichtenthaler H.K. (1987). Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. - *Methods in Enzymology*, 148: 350-382.
- McCune D. C., Hitchcock A. E. and Jacobson J. S. (1965). Fluoride accumulation and growth of plants exposed to particulate cryolite in the atmosphere. *Contributions from Boyce Thompson Institute*, 23: 1-22.
- Miller R.J., Bittell J. E. and Koeppel D. E. (1973). The effect of cadmium on electron and energy transfer reactions in corn mitochondria, *Physiologia Plantarum* 28: 166-171.
- Miller P. M. and Rick S. (1967). Shoot damage to green house plants. *Plant Disease Reporter*, 51: 712.

- Moraga S. P., Escobar R. and Valenzuela A. S. (2006). Resistance to freezing in three *Eucalyptus globules* Labill subspecies. *Electronic Journal of Biotechnology*, 9:310-314.
- Naya L., Ladrera R., Ramos J., Gonzalez E. M., Arreseigor C., Minchin F. R. and Becana M. (2007). The response of carbon metabolism and antioxidant defenses of alfalfa nodules to drought stress and to the subsequent recovery of plants. *Plant Physiology*, 114:1104-1114.
- Pandey D. D. and Simba A. K. (1989). Effect of cement kiln dust on chlorophyll in gram leaves. *Indian Journal of Ecology*, 16: 178-179.
- Pandey D. D. and Simba A. K. (1988). Effect of cement dust pollution on growth parameters of maize crop. *Proceeding of Botany Section of 75th India Science Congress*. 313, Pune, India.
- Pandey D. D. and Simba A. K. (1990). Effect of cement kiln dust on chlorophyll in wheat leaf. *Environment and Ecology*, 8: 461-463.
- Pandey D. D., Nirala A. K. and Gaulam R. R. (1999). Impact of stone crusher dust pollution on maize crop. *Indian Journal of Environment and Ecoplanning*, 2: 43-46.
- Parthasarathy S., Arunachalam M. and Natarajan K. (1975). Effect of cement dust pollution on certain physical parameter of maize crop and soils. *Indian Journal of Environmental Health*, 17: 114-120.
- Rao D. N. (1971). Study of the air pollution problem due to coal unloading in Varanasi, India. *Proceedings of the 2nd Inter Clean Air Congress* (Engly HM, Berry WT, eds).pp. 273-276, Academic Press, New York, USA
- Santosh K.P. and Tripathi B. D. (2008). Seasonal Variation of Leaf Dust Accumulation and Pigment Content in Plant Species Exposed to Urban Particulates Pollution. *Journal of Environmental Quality*, 37:865-870.
- Schutzki R.E. and Cregg B. (2007). *Abiotic Plant Disorders Symptoms, Signs and Solutions. A Diagnostic Guide to Problem Solving*. Extension Bulletin. E-2996.
- Seyyednejad S.M., Niknejad M. and Yusefi M. (2009). Study of air pollution effects on some physiology and morphology factors of *Albizia lebbek* in high temperature condition in Khuzestan. *Journal of Plant Sciences*, 4: 122-126
- Seyyednejad S.M., Niknejad M. and Koochak H. (2011). A review of some different effects of air pollution on plants. *Research Journal of Environmental Sciences*, 5:302-309
- Seyyednejad S.M., Majdian K., Koochak H. and Niknejad M. (2011). Air pollution tolerance indices of some plants around industrial zone in south of Iran. *Asian Journal of Biological Sciences*, 4: 300-305
- Shweta M. and Agrawal S. B. (2006). Interactive effects between supplemental ultraviolet-B and heavy metals on the growth and biochemical characteristics of *Spinacia oleracea* L. *Brazil. J. Plant Physiology*, 18: 98-102.

- Shvaleva A. L., Costa F., Silva E., Breia E., Jouve L., Hausman J.F., Almeida M. H., Maroco J. P., Rodrigues M. L., Pereira J.S. and Chaves M. M. (2005). Metabolic responses to water deficit in two Eucalyptus globules clones with contrasting drought sensitivity. *Tree Physiology*, 26:239-248.
- Singh S. N. and Rao D. N. (1978). Effect of cement dust pollution on soil properties and on wheat plants. *Indian Journal of Environmental Health*, 20: 258-267.
- Stern A. C. (1976). *Air Pollution, Measurement, Monitoring and surveillance of air pollution*, 3rd ed., Academic press, Newyork.
- Taylor G. and Davies W. J. (1990). Root growth of *Fagus sylvatica*: Impact of air quality and drought at a site in southern Britain. *New Phytologist*. 166:457-464.
- Thayumanavan B. and Sadasivam S. (1984). Carbohydrate chemistry. *Quality of Plant Foods Human Nutrition*, 34: 253-254.
- Thambavani S. D. and kumar S. R. (2011). Changes of Photosynthetic pigments in some selected Plant species induced by cement dust pollution. *The Ecoscan*, 1: 167-172.
- Thambavani S. D. and kumar S. R. (2011). The monthly changes of chloroplast pigments content in selected plant species exposed to cement dust pollution. *Journal of research in Biology*. 8: 660-666.
- Thambavani S. D. and kumar S. R. (2012). Correlation Analysis and Exceedence Factor among the Ambient Gaseous Pollutants and Particulate Matter in Urban Area. *Journal of research in Biology*, 3: 232-240.
- Tiwari, S., M. Agrawal and F.M. Marshall. (2006). Evaluation of ambient air pollution impact on carrot plants at a sub urban site using open top chambers. *Environment Monitoring Assessment*, 119: 15-30.
- Tripathi A. K. and Mukesh G. (2006). Biochemical parameters of plants as indicators of air pollution. *Journal of Environmental Biology*, 28:127-132.
- Tripathi A. K. and Gautam M. (2007). Biochemical parameters of plants as indicators of air pollution. *Journal of Environmental Biology*, 28: 127-132.

**تأثير تلوث الهواء على الصفات المورفولوجية والفسيزيولوجية في نبات الخبيزة
محمد السيد أبو زيادة، سامية على هارون، غادة عبدالله الشربيني و أحمد عبد الله نجم
قسم النبات - كلية العلوم - جامعة المنصورة - مصر**

زيادة الأنشطة الصناعية أدى إلى ظهور العديد من ملوثات الهواء في الغلاف الجوي هذه الملوثات لها تأثير خطير على الإنسان والنبات. هذه الدراسة تصنف تأثير ملوثات الهواء الناتجة من مصانع الطوب و طرق المواصلات على الخصائص المورفولوجية و الفسيزيولوجية لنبات الخبيزة التي تنمو طبيعيا حول هذه المصادر . تم قياس الخصائص المورفولوجية مثل طول الساق & عدد الافرع وطول الجذر وطول وعرض الورقة ومساحة الورقة بالإضافة لبعض الخصائص الفسيزيولوجية مثل مستوى كلوروفيل (أ & ب) والكاروتينات والسكريات الكلية في النبات حيث اظهرت النتائج انخفاض ملحوظ في هذه الخصائص في النباتات القريبة من مصادر التلوث .